

表 3775 Non-HDL-C濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	183	846	1,870	2,408	2,009	1,281	624	270	93	42	9,626	5,450	15,076	108.12	32.83
(%)	(1.9)	(8.8)	(19.4)	(25.0)	(20.9)	(13.3)	(6.5)	(2.8)	(1.0)	(0.4)	(100.0)				
慢性腎盂腎炎	3	18	56	86	58	38	17	10	2	1	289	133	422	109.44	31.51
(%)	(1.0)	(6.2)	(19.4)	(29.8)	(20.1)	(13.1)	(5.9)	(3.5)	(0.7)	(0.3)	(100.0)				
急速進行性糸球体腎炎	2	12	33	55	37	24	16	5	2	2	188	96	284	111.47	34.79
(%)	(1.1)	(6.4)	(17.6)	(29.3)	(19.7)	(12.8)	(8.5)	(2.7)	(1.1)	(1.1)	(100.0)				
妊娠腎/妊娠中毒症	3	9	38	60	55	46	23	12	5		251	108	359	117.37	32.73
(%)	(1.2)	(3.6)	(15.1)	(23.9)	(21.9)	(18.3)	(9.2)	(4.8)	(2.0)		(100.0)				
その他の分類不能の腎炎	3	13	23	30	23	21	8	3	1		125	70	195	106.78	31.98
(%)	(2.4)	(10.4)	(18.4)	(24.0)	(18.4)	(16.8)	(6.4)	(2.4)	(0.8)		(100.0)				
多発性嚢胞腎	13	93	193	241	204	105	58	22	5	3	937	527	1,464	105.86	30.86
(%)	(1.4)	(9.9)	(20.6)	(25.7)	(21.8)	(11.2)	(6.2)	(2.3)	(0.5)	(0.3)	(100.0)				
腎硬化症	38	152	397	461	376	234	112	40	17	15	1,842	1,044	2,886	107.36	33.15
(%)	(2.1)	(8.3)	(21.6)	(25.0)	(20.4)	(12.7)	(6.1)	(2.2)	(0.9)	(0.8)	(100.0)				
悪性高血圧	1	13	38	42	54	27	11	8	2	2	198	114	312	112.30	35.68
(%)	(0.5)	(6.6)	(19.2)	(21.2)	(27.3)	(13.6)	(5.6)	(4.0)	(1.0)	(1.0)	(100.0)				
糖尿病性腎症	203	925	1,772	1,985	1,631	922	487	226	92	45	8,288	4,885	13,173	105.09	33.87
(%)	(2.4)	(11.2)	(21.4)	(24.0)	(19.7)	(11.1)	(5.9)	(2.7)	(1.1)	(0.5)	(100.0)				
SLE腎炎	8	16	47	41	48	33	24	6	5	5	233	130	363	113.56	39.13
(%)	(3.4)	(6.9)	(20.2)	(17.6)	(20.6)	(14.2)	(10.3)	(2.6)	(2.1)	(2.1)	(100.0)				
アミロイド腎	2	5	4	9	9	4		4	1		38	21	59	109.03	39.94
(%)	(5.3)	(13.2)	(10.5)	(23.7)	(23.7)	(10.5)		(10.5)	(2.6)		(100.0)				
痛風腎		15	25	20	16	12	4	4	2	1	99	55	154	106.03	35.66
(%)		(15.2)	(25.3)	(20.2)	(16.2)	(12.1)	(4.0)	(4.0)	(2.0)	(1.0)	(100.0)				
先天性代謝異常による腎不全		5	2	9	3	5	2		1		27	17	44	108.11	34.76
(%)		(18.5)	(7.4)	(33.3)	(11.1)	(18.5)	(7.4)		(3.7)		(100.0)				
腎・尿路結核		3	2	3	4		2				14	8	22	101.57	32.33
(%)		(21.4)	(14.3)	(21.4)	(28.6)		(14.3)				(100.0)				
腎・尿路結石		5	14	8	4	3	2		1		37	32	69	98.54	32.04
(%)		(13.5)	(37.8)	(21.6)	(10.8)	(8.1)	(5.4)		(2.7)		(100.0)				
腎・尿路腫瘍	3	4	18	18	10	5	4	2			64	21	85	100.83	30.85
(%)	(4.7)	(6.3)	(28.1)	(28.1)	(15.6)	(7.8)	(6.3)	(3.1)			(100.0)				
閉塞性尿路障害		1	10	13	8	5	3	1	2		43	31	74	111.72	33.56
(%)		(2.3)	(23.3)	(30.2)	(18.6)	(11.6)	(7.0)	(2.3)	(4.7)		(100.0)				
骨髄腫	1	2	2	6	3				1		15	4	19	97.80	35.64
(%)	(6.7)	(13.3)	(13.3)	(40.0)	(20.0)				(6.7)		(100.0)				
腎形成不全	1	6	13	16	17	7	3	2		1	66	35	101	107.83	32.72
(%)	(1.5)	(9.1)	(19.7)	(24.2)	(25.8)	(10.6)	(4.5)	(3.0)		(1.5)	(100.0)				
不明	45	183	426	533	421	272	135	53	19	18	2,105	1,375	3,480	107.80	33.45
(%)	(2.1)	(8.7)	(20.2)	(25.3)	(20.0)	(12.9)	(6.4)	(2.5)	(0.9)	(0.9)	(100.0)				
再導入	8	22	44	76	58	29	22	4	2	4	269	147	416	109.39	36.90
(%)	(3.0)	(8.2)	(16.4)	(28.3)	(21.6)	(10.8)	(8.2)	(1.5)	(0.7)	(1.5)	(100.0)				
その他	12	55	111	145	114	76	40	10	9	2	574	315	889	107.18	32.81
(%)	(2.1)	(9.6)	(19.3)	(25.3)	(19.9)	(13.2)	(7.0)	(1.7)	(1.6)	(0.3)	(100.0)				
合計	529	2,403	5,138	6,265	5,162	3,149	1,597	682	262	141	25,328	14,618	39,946	107.12	33.37
(%)	(2.1)	(9.5)	(20.3)	(24.7)	(20.4)	(12.4)	(6.3)	(2.7)	(1.0)	(0.6)	(100.0)				
記載なし															
(%)															
総計	529	2,403	5,138	6,265	5,162	3,149	1,597	682	262	141	25,328	14,618	39,946	107.12	33.37
(%)	(2.1)	(9.5)	(20.3)	(24.7)	(20.4)	(12.4)	(6.3)	(2.7)	(1.0)	(0.6)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査