

表 3792 透析前収縮期血圧(mmHg) 併用の状況別（透析患者全体）

併用の状況	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	39	419	3,209	16,946	51,681	79,145	57,424	21,983	5,509	1,148	237,503	63,291	300,794	151.34	24.21
(%)	(0.0)	(0.2)	(1.4)	(7.1)	(21.8)	(33.3)	(24.2)	(9.3)	(2.3)	(0.5)	(100.0)				
PD単独	1	13	100	517	1,234	1,136	443	114	13	3	3,574	3,614	7,188	138.34	21.77
(%)	(0.0)	(0.4)	(2.8)	(14.5)	(34.5)	(31.8)	(12.4)	(3.2)	(0.4)	(0.1)	(100.0)				
HD(F)単独.PDカテあり		1	3	12	53	68	35	15	5	2	194	75	269	149.73	25.58
(%)		(0.5)	(1.5)	(6.2)	(27.3)	(35.1)	(18.0)	(7.7)	(2.6)	(1.0)	(100.0)				
PD.HD併用週1 HD(F)			21	113	250	386	247	88	16	7	1,128	416	1,544	148.34	24.38
(%)			(1.9)	(10.0)	(22.2)	(34.2)	(21.9)	(7.8)	(1.4)	(0.6)	(100.0)				
PD.HD併用週2 HD(F)			3	9	32	45	26	16	2		133	44	177	149.88	24.16
(%)			(2.3)	(6.8)	(24.1)	(33.8)	(19.5)	(12.0)	(1.5)		(100.0)				
PD.HD併用週3 HD(F)				4	5	13	6	3			31	9	40	147.10	22.87
(%)				(12.9)	(16.1)	(41.9)	(19.4)	(9.7)			(100.0)				
PD.HD併用週4 HD(F)					1						1		1	130.00	
(%)					(100.0)						(100.0)				
PD.HD併用その他			1	5	12	23	12	4			57	16	73	147.28	20.48
(%)			(1.8)	(8.8)	(21.1)	(40.4)	(21.1)	(7.0)			(100.0)				
合計	40	433	3,337	17,606	53,268	80,816	58,193	22,223	5,545	1,160	242,621	67,465	310,086	151.13	24.23
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.3)	(24.0)	(9.2)	(2.3)	(0.5)	(100.0)				
不明			1	1	3	6	2	1	1		15	7	22	147.13	30.55
(%)			(6.7)	(6.7)	(20.0)	(40.0)	(13.3)	(6.7)	(6.7)		(100.0)				
記載なし (%)															
総計	40	433	3,338	17,607	53,271	80,822	58,195	22,224	5,546	1,160	242,636	67,472	310,108	151.13	24.23
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.3)	(24.0)	(9.2)	(2.3)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査