

表 3793 透析前収縮期血圧(mmHg) 性別 (透析患者全体)

性別	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
男性	15	224	1,803	10,335	33,344	52,546	38,126	14,493	3,505	693	155,084	43,057	198,141	151.76	23.73
(%)	(0.0)	(0.1)	(1.2)	(6.7)	(21.5)	(33.9)	(24.6)	(9.3)	(2.3)	(0.4)	(100.0)				
女性	25	209	1,535	7,272	19,927	28,276	20,069	7,731	2,041	467	87,552	24,415	111,967	150.01	25.05
(%)	(0.0)	(0.2)	(1.8)	(8.3)	(22.8)	(32.3)	(22.9)	(8.8)	(2.3)	(0.5)	(100.0)				
合計	40	433	3,338	17,607	53,271	80,822	58,195	22,224	5,546	1,160	242,636	67,472	310,108	151.13	24.23
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.3)	(24.0)	(9.2)	(2.3)	(0.5)	(100.0)				
記載なし															
(%)															
総計	40	433	3,338	17,607	53,271	80,822	58,195	22,224	5,546	1,160	242,636	67,472	310,108	151.13	24.23
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.3)	(24.0)	(9.2)	(2.3)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査