

表 3797 透析前収縮期血圧(mmHg) 医療機関別（透析患者全体）

医療機関	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
国公立大学		3	34	134	264	268	126	38	6	3	876	381	1,257	139.16	23.68
(%)		(0.3)	(3.9)	(15.3)	(30.1)	(30.6)	(14.4)	(4.3)	(0.7)	(0.3)	(100.0)				
私立大学		9	44	229	604	736	477	177	41	11	2,328	1,189	3,517	147.04	24.75
(%)		(0.4)	(1.9)	(9.8)	(25.9)	(31.6)	(20.5)	(7.6)	(1.8)	(0.5)	(100.0)				
国立		3	17	48	141	203	115	52	11	1	591	278	869	147.69	24.89
(%)		(0.5)	(2.9)	(8.1)	(23.9)	(34.3)	(19.5)	(8.8)	(1.9)	(0.2)	(100.0)				
県市町村立	3	25	234	1,121	3,203	4,595	3,338	1,245	334	61	14,159	5,026	19,185	150.39	24.63
(%)	(0.0)	(0.2)	(1.7)	(7.9)	(22.6)	(32.5)	(23.6)	(8.8)	(2.4)	(0.4)	(100.0)				
社会保険	1	8	50	236	794	1,073	784	266	51	10	3,273	599	3,872	149.69	23.46
(%)	(0.0)	(0.2)	(1.5)	(7.2)	(24.3)	(32.8)	(24.0)	(8.1)	(1.6)	(0.3)	(100.0)				
厚生連		10	96	571	1,723	2,440	1,847	729	190	35	7,641	1,623	9,264	151.51	24.49
(%)		(0.1)	(1.3)	(7.5)	(22.5)	(31.9)	(24.2)	(9.5)	(2.5)	(0.5)	(100.0)				
その他公的	1	28	107	692	1,862	2,816	1,882	699	178	38	8,303	2,686	10,989	149.84	24.39
(%)	(0.0)	(0.3)	(1.3)	(8.3)	(22.4)	(33.9)	(22.7)	(8.4)	(2.1)	(0.5)	(100.0)				
私立総合		9	74	477	1,327	1,959	1,414	563	148	43	6,014	1,540	7,554	151.36	24.81
(%)		(0.1)	(1.2)	(7.9)	(22.1)	(32.6)	(23.5)	(9.4)	(2.5)	(0.7)	(100.0)				
私立	15	137	1,098	5,361	15,193	21,908	15,737	6,151	1,502	315	67,417	21,225	88,642	150.39	24.62
(%)	(0.0)	(0.2)	(1.6)	(8.0)	(22.5)	(32.5)	(23.3)	(9.1)	(2.2)	(0.5)	(100.0)				
私立診療所	20	201	1,584	8,738	28,160	44,824	32,475	12,304	3,085	643	132,034	32,925	164,959	151.83	23.89
(%)	(0.0)	(0.2)	(1.2)	(6.6)	(21.3)	(33.9)	(24.6)	(9.3)	(2.3)	(0.5)	(100.0)				
総計	40	433	3,338	17,607	53,271	80,822	58,195	22,224	5,546	1,160	242,636	67,472	310,108	151.13	24.23
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.3)	(24.0)	(9.2)	(2.3)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査