

表 3798 透析前収縮期血圧(mmHg) 週透析回数(回/週)別（透析患者全体）

週透析回数	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
1回		1	34	158	372	548	346	127	22	7	1,615	225	1,840	147.84	24.24
(%)		(0.1)	(2.1)	(9.8)	(23.0)	(33.9)	(21.4)	(7.9)	(1.4)	(0.4)	(100.0)				
2回	1	14	79	443	1,244	1,778	1,166	413	86	21	5,245	947	6,192	148.86	23.92
(%)	(0.0)	(0.3)	(1.5)	(8.4)	(23.7)	(33.9)	(22.2)	(7.9)	(1.6)	(0.4)	(100.0)				
3回	38	392	3,047	16,146	49,583	76,109	55,424	21,235	5,349	1,112	228,435	42,204	270,639	151.44	24.19
(%)	(0.0)	(0.2)	(1.3)	(7.1)	(21.7)	(33.3)	(24.3)	(9.3)	(2.3)	(0.5)	(100.0)				
4回		6	33	103	212	277	156	64	19	3	873	121	994	144.84	26.89
(%)		(0.7)	(3.8)	(11.8)	(24.3)	(31.7)	(17.9)	(7.3)	(2.2)	(0.3)	(100.0)				
5回			2	15	17	19	12	1	1		67	22	89	139.15	24.97
(%)			(3.0)	(22.4)	(25.4)	(28.4)	(17.9)	(1.5)	(1.5)		(100.0)				
6回			3	11	17	13	10	2	1		57	24	81	137.74	23.98
(%)			(5.3)	(19.3)	(29.8)	(22.8)	(17.5)	(3.5)	(1.8)		(100.0)				
7回		1	2	5	3	7	3				21	9	30	130.62	26.83
(%)		(4.8)	(9.5)	(23.8)	(14.3)	(33.3)	(14.3)				(100.0)				
合計	39	414	3,200	16,881	51,448	78,751	57,117	21,842	5,478	1,143	236,313	43,552	279,865	151.32	24.21
(%)	(0.0)	(0.2)	(1.4)	(7.1)	(21.8)	(33.3)	(24.2)	(9.2)	(2.3)	(0.5)	(100.0)				
記載なし	1	19	138	726	1,823	2,071	1,078	382	68	17	6,323	23,920	30,243	143.91	23.80
(%)	(0.0)	(0.3)	(2.2)	(11.5)	(28.8)	(32.8)	(17.0)	(6.0)	(1.1)	(0.3)	(100.0)				
総計	40	433	3,338	17,607	53,271	80,822	58,195	22,224	5,546	1,160	242,636	67,472	310,108	151.13	24.23
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.3)	(24.0)	(9.2)	(2.3)	(0.5)	(100.0)				
平均	2.97	2.99	2.97	2.97	2.97	2.97	2.97	2.97	2.98	2.97	2.97	2.97	2.97		
標準偏差	0.16	0.31	0.31	0.29	0.25	0.24	0.22	0.21	0.19	0.21	0.24	0.23	0.24		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査