

表 3803 透析前収縮期血圧(mmHg) 透析歴(年)別 (施設血液透析患者 透析回数週3回)

透析歴	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
2年未満	9	38	454	3,111	9,817	14,298	9,896	3,784	912	171	42,490	8,225	50,715	150.75	23.67
(%)	(0.0)	(0.1)	(1.1)	(7.3)	(23.1)	(33.7)	(23.3)	(8.9)	(2.1)	(0.4)	(100.0)				
2年～	7	50	477	2,910	10,067	16,380	12,464	4,975	1,291	274	48,895	9,394	58,289	153.16	23.83
(%)	(0.0)	(0.1)	(1.0)	(6.0)	(20.6)	(33.5)	(25.5)	(10.2)	(2.6)	(0.6)	(100.0)				
5年～	5	50	496	2,928	9,798	16,555	12,960	5,214	1,400	271	49,677	9,550	59,227	153.73	23.99
(%)	(0.0)	(0.1)	(1.0)	(5.9)	(19.7)	(33.3)	(26.1)	(10.5)	(2.8)	(0.5)	(100.0)				
10年～	3	32	295	1,676	5,467	8,476	6,273	2,345	577	103	25,247	4,785	30,032	151.74	23.72
(%)	(0.0)	(0.1)	(1.2)	(6.6)	(21.7)	(33.6)	(24.8)	(9.3)	(2.3)	(0.4)	(100.0)				
15年～	4	31	227	979	3,005	4,408	2,856	989	193	55	12,747	2,502	15,249	148.96	23.76
(%)	(0.0)	(0.2)	(1.8)	(7.7)	(23.6)	(34.6)	(22.4)	(7.8)	(1.5)	(0.4)	(100.0)				
20年～	1	14	159	635	1,712	2,221	1,369	393	87	17	6,608	1,323	7,931	145.99	23.61
(%)	(0.0)	(0.2)	(2.4)	(9.6)	(25.9)	(33.6)	(20.7)	(5.9)	(1.3)	(0.3)	(100.0)				
25年～	2	19	108	450	925	1,027	594	189	33	2	3,349	695	4,044	142.18	24.62
(%)	(0.1)	(0.6)	(3.2)	(13.4)	(27.6)	(30.7)	(17.7)	(5.6)	(1.0)	(0.1)	(100.0)				
30年～	2	14	85	251	494	564	289	72	12		1,783	400	2,183	139.43	24.59
(%)	(0.1)	(0.8)	(4.8)	(14.1)	(27.7)	(31.6)	(16.2)	(4.0)	(0.7)		(100.0)				
35年～	1	14	59	166	231	232	118	37	10	1	869	164	1,033	136.48	26.92
(%)	(0.1)	(1.6)	(6.8)	(19.1)	(26.6)	(26.7)	(13.6)	(4.3)	(1.2)	(0.1)	(100.0)				
40年～	1	9	17	48	51	35	19	4			184	37	221	126.82	27.17
(%)	(0.5)	(4.9)	(9.2)	(26.1)	(27.7)	(19.0)	(10.3)	(2.2)			(100.0)				
合計	35	271	2,377	13,154	41,567	64,196	46,838	18,002	4,515	894	191,849	37,075	228,924	151.64	24.02
(%)	(0.0)	(0.1)	(1.2)	(6.9)	(21.7)	(33.5)	(24.4)	(9.4)	(2.4)	(0.5)	(100.0)				
不明				1	9	7	5				22	25	47	145.36	17.51
(%)				(4.5)	(40.9)	(31.8)	(22.7)				(100.0)				
記載なし															
(%)															
総計	35	271	2,377	13,155	41,576	64,203	46,843	18,002	4,515	894	191,871	37,100	228,971	151.64	24.02
(%)	(0.0)	(0.1)	(1.2)	(6.9)	(21.7)	(33.5)	(24.4)	(9.4)	(2.4)	(0.5)	(100.0)				
平均	10.83	13.17	10.27	8.19	7.28	6.89	6.61	6.25	5.98	5.97	6.96	7.04	6.98		
標準偏差	11.79	12.04	10.03	8.64	7.52	6.88	6.39	5.95	5.59	5.24	7.02	7.14	7.04		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査