

表 3805 透析前収縮期血圧(mmHg) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
国公立大学			14	38	85	120	71	23	6	3	360	54	414	145.20	25.48
(%)			(3.9)	(10.6)	(23.6)	(33.3)	(19.7)	(6.4)	(1.7)	(0.8)	(100.0)				
私立大学		4	27	138	323	447	328	134	36	9	1,446	122	1,568	149.82	25.54
(%)		(0.3)	(1.9)	(9.5)	(22.3)	(30.9)	(22.7)	(9.3)	(2.5)	(0.6)	(100.0)				
国立		2	12	23	88	146	100	43	11	1	426	69	495	151.12	24.83
(%)		(0.5)	(2.8)	(5.4)	(20.7)	(34.3)	(23.5)	(10.1)	(2.6)	(0.2)	(100.0)				
県市町村立	3	14	156	772	2,353	3,469	2,638	988	263	47	10,703	2,127	12,830	151.39	24.35
(%)	(0.0)	(0.1)	(1.5)	(7.2)	(22.0)	(32.4)	(24.6)	(9.2)	(2.5)	(0.4)	(100.0)				
社会保険		2	33	172	608	863	671	222	45	7	2,623	173	2,796	151.05	22.88
(%)		(0.1)	(1.3)	(6.6)	(23.2)	(32.9)	(25.6)	(8.5)	(1.7)	(0.3)	(100.0)				
厚生連		6	65	424	1,336	1,943	1,487	612	160	28	6,061	527	6,588	152.39	24.36
(%)		(0.1)	(1.1)	(7.0)	(22.0)	(32.1)	(24.5)	(10.1)	(2.6)	(0.5)	(100.0)				
その他公的	1	20	66	478	1,429	2,222	1,569	568	153	27	6,533	732	7,265	150.97	24.00
(%)	(0.0)	(0.3)	(1.0)	(7.3)	(21.9)	(34.0)	(24.0)	(8.7)	(2.3)	(0.4)	(100.0)				
私立総合		6	51	349	952	1,480	1,078	428	124	29	4,497	487	4,984	151.97	24.77
(%)		(0.1)	(1.1)	(7.8)	(21.2)	(32.9)	(24.0)	(9.5)	(2.8)	(0.6)	(100.0)				
私立	13	93	835	4,197	12,318	17,938	12,945	5,139	1,277	256	55,011	11,588	66,599	150.81	24.52
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.4)	(32.6)	(23.5)	(9.3)	(2.3)	(0.5)	(100.0)				
私立診療所	18	124	1,118	6,564	22,084	35,575	25,956	9,845	2,440	487	104,211	21,221	125,432	152.16	23.64
(%)	(0.0)	(0.1)	(1.1)	(6.3)	(21.2)	(34.1)	(24.9)	(9.4)	(2.3)	(0.5)	(100.0)				
総計	35	271	2,377	13,155	41,576	64,203	46,843	18,002	4,515	894	191,871	37,100	228,971	151.64	24.02
(%)	(0.0)	(0.1)	(1.2)	(6.9)	(21.7)	(33.5)	(24.4)	(9.4)	(2.4)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査