

表 3815 透析前収縮期血圧(mmHg) 透析前拡張期血圧(mmHg)別（血液透析濾過患者 透析回数週3回）

透析前拡張 期血圧	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
20未満			1			1	3	2			7		7	157.43	38.11
(%)			(14.3)			(14.3)	(42.9)	(28.6)			(100.0)				
20～	1	27	26	36	18	8	3	1	1		121		121	104.28	28.34
(%)	(0.8)	(22.3)	(21.5)	(29.8)	(14.9)	(6.6)	(2.5)	(0.8)	(0.8)		(100.0)				
40～	1	84	427	888	770	393	91	22	2	1	2,679	1	2,680	119.54	22.85
(%)	(0.0)	(3.1)	(15.9)	(33.1)	(28.7)	(14.7)	(3.4)	(0.8)	(0.1)	(0.0)	(100.0)				
60～		4	155	1,725	5,049	5,315	2,247	420	50	7	14,972		14,972	141.61	19.62
(%)		(0.0)	(1.0)	(11.5)	(33.7)	(35.5)	(15.0)	(2.8)	(0.3)	(0.0)	(100.0)				
80～			1	129	1,457	3,723	2,756	800	120	17	9,003		9,003	156.05	18.20
(%)			(0.0)	(1.4)	(16.2)	(41.4)	(30.6)	(8.9)	(1.3)	(0.2)	(100.0)				
90～				10	269	1,599	2,181	984	238	36	5,317		5,317	167.09	18.52
(%)				(0.2)	(5.1)	(30.1)	(41.0)	(18.5)	(4.5)	(0.7)	(100.0)				
100～				3	42	348	971	868	362	107	2,701		2,701	180.40	20.46
(%)				(0.1)	(1.6)	(12.9)	(35.9)	(32.1)	(13.4)	(4.0)	(100.0)				
120～						7	27	43	48	44	169		169	200.96	24.24
(%)						(4.1)	(16.0)	(25.4)	(28.4)	(26.0)	(100.0)				
140～							2	4	1	4	11	1	12	207.09	31.13
(%)							(18.2)	(36.4)	(9.1)	(36.4)	(100.0)				
合計	2	115	610	2,791	7,605	11,394	8,281	3,144	822	216	34,980	2	34,982	150.69	25.04
(%)	(0.0)	(0.3)	(1.7)	(8.0)	(21.7)	(32.6)	(23.7)	(9.0)	(2.3)	(0.6)	(100.0)				
記載なし			5	9	34	39	27	10			124	4,840	4,964	144.16	23.77
(%)			(4.0)	(7.3)	(27.4)	(31.5)	(21.8)	(8.1)			(100.0)				
総計	2	115	615	2,800	7,639	11,433	8,308	3,154	822	216	35,104	4,842	39,946	150.67	25.04
(%)	(0.0)	(0.3)	(1.8)	(8.0)	(21.8)	(32.6)	(23.7)	(9.0)	(2.3)	(0.6)	(100.0)				
平均	36.00	44.12	53.50	63.33	71.81	79.08	85.95	92.46	99.63	108.05	79.17	121.00	79.17		
標準偏差	9.90	7.64	8.57	9.76	10.37	11.06	11.92	12.89	13.45	16.28	14.69	87.68	14.70		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査