

表 3816 透析前収縮期血圧(mmHg) 透析前脈拍(拍/分)別（血液透析濾過患者 透析回数週3回）

透析前脈拍	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
40未満			1	2	6	3	2	2		1	17		17	142.59	34.39
(%)			(5.9)	(11.8)	(35.3)	(17.6)	(11.8)	(11.8)		(5.9)	(100.0)				
40～		6	7	40	65	93	62	18	4		295		295	144.46	26.40
(%)		(2.0)	(2.4)	(13.6)	(22.0)	(31.5)	(21.0)	(6.1)	(1.4)		(100.0)				
50～		8	58	249	674	884	539	171	40	9	2,632		2,632	146.76	23.98
(%)		(0.3)	(2.2)	(9.5)	(25.6)	(33.6)	(20.5)	(6.5)	(1.5)	(0.3)	(100.0)				
60～		11	139	684	2,129	2,855	1,835	638	158	37	8,486	4	8,490	148.68	23.74
(%)		(0.1)	(1.6)	(8.1)	(25.1)	(33.6)	(21.6)	(7.5)	(1.9)	(0.4)	(100.0)				
70～	1	33	162	799	2,239	3,501	2,576	952	227	54	10,544	1	10,545	151.03	24.47
(%)	(0.0)	(0.3)	(1.5)	(7.6)	(21.2)	(33.2)	(24.4)	(9.0)	(2.2)	(0.5)	(100.0)				
80～	1	21	104	508	1,344	2,254	1,811	733	228	55	7,059	4	7,063	153.46	25.62
(%)	(0.0)	(0.3)	(1.5)	(7.2)	(19.0)	(31.9)	(25.7)	(10.4)	(3.2)	(0.8)	(100.0)				
90～		16	65	251	618	954	857	379	113	39	3,292	1	3,293	153.97	27.07
(%)		(0.5)	(2.0)	(7.6)	(18.8)	(29.0)	(26.0)	(11.5)	(3.4)	(1.2)	(100.0)				
100～		4	30	115	181	291	230	142	35	14	1,042		1,042	152.26	28.96
(%)		(0.4)	(2.9)	(11.0)	(17.4)	(27.9)	(22.1)	(13.6)	(3.4)	(1.3)	(100.0)				
110～		5	9	30	36	83	64	27	4	3	261		261	149.10	28.69
(%)		(1.9)	(3.4)	(11.5)	(13.8)	(31.8)	(24.5)	(10.3)	(1.5)	(1.1)	(100.0)				
120～		3	8	15	16	25	29	6	4		106		106	144.48	31.15
(%)		(2.8)	(7.5)	(14.2)	(15.1)	(23.6)	(27.4)	(5.7)	(3.8)		(100.0)				
合計	2	107	583	2,693	7,308	10,943	8,005	3,068	813	212	33,734	10	33,744	150.84	25.10
(%)	(0.0)	(0.3)	(1.7)	(8.0)	(21.7)	(32.4)	(23.7)	(9.1)	(2.4)	(0.6)	(100.0)				
記載なし		8	32	107	331	490	303	86	9	4	1,370	4,832	6,202	146.35	23.10
(%)		(0.6)	(2.3)	(7.8)	(24.2)	(35.8)	(22.1)	(6.3)	(0.7)	(0.3)	(100.0)				
総計	2	115	615	2,800	7,639	11,433	8,308	3,154	822	216	35,104	4,842	39,946	150.67	25.04
(%)	(0.0)	(0.3)	(1.8)	(8.0)	(21.8)	(32.6)	(23.7)	(9.0)	(2.3)	(0.6)	(100.0)				
平均	76.50	79.39	76.36	75.27	73.85	74.85	76.26	77.69	78.67	80.07	75.43	77.80	75.43		
標準偏差	6.36	17.53	15.43	14.21	12.59	12.55	12.75	12.89	13.20	13.08	12.93	10.40	12.93		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査