

表 3826 透析前拡張期血圧(mmHg) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
施設血液透析	71	646	17,149	93,525	48,769	25,897	12,580	771	62	199,470	56,171	255,641	77.52	14.39
(%)	(0.0)	(0.3)	(8.6)	(46.9)	(24.4)	(13.0)	(6.3)	(0.4)	(0.0)	(100.0)				
Off-line HDF	1	38	543	2,342	1,203	603	312	30	2	5,074	1,241	6,315	76.34	15.29
(%)	(0.0)	(0.7)	(10.7)	(46.2)	(23.7)	(11.9)	(6.1)	(0.6)	(0.0)	(100.0)				
On-line HDF	5	83	2,134	12,745	7,859	4,774	2,410	142	10	30,162	5,928	36,090	79.69	14.53
(%)	(0.0)	(0.3)	(7.1)	(42.3)	(26.1)	(15.8)	(8.0)	(0.5)	(0.0)	(100.0)				
Push/Pull HDF		1	43	201	119	45	17			426	111	537	76.41	13.04
(%)		(0.2)	(10.1)	(47.2)	(27.9)	(10.6)	(4.0)			(100.0)				
ハイオフィльтраーション	1	2	44	121	61	40	18			287	54	341	75.00	16.76
(%)	(0.3)	(0.7)	(15.3)	(42.2)	(21.3)	(13.9)	(6.3)			(100.0)				
血液濾過			3	44	22	14	5			88	4	92	78.88	12.12
(%)			(3.4)	(50.0)	(25.0)	(15.9)	(5.7)			(100.0)				
血液吸着透析 (リクセル)	1	10	149	712	272	118	53	3		1,318	305	1,623	73.92	14.13
(%)	(0.1)	(0.8)	(11.3)	(54.0)	(20.6)	(9.0)	(4.0)	(0.2)		(100.0)				
血液吸着透析 (その他)			2	2	1					5	2	7	67.20	15.16
(%)			(40.0)	(40.0)	(20.0)					(100.0)				
在宅血液透析		1	38	121	85	46	29			320	201	521	78.73	15.04
(%)		(0.3)	(11.9)	(37.8)	(26.6)	(14.4)	(9.1)			(100.0)				
腹膜透析 (CAPD)	1	8	213	1,140	667	346	171	10		2,556	2,465	5,021	78.26	14.40
(%)	(0.0)	(0.3)	(8.3)	(44.6)	(26.1)	(13.5)	(6.7)	(0.4)		(100.0)				
腹膜透析 (APD)		6	115	662	432	233	125	4		1,577	1,262	2,839	79.47	14.29
(%)		(0.4)	(7.3)	(42.0)	(27.4)	(14.8)	(7.9)	(0.3)		(100.0)				
腹膜透析 (CCPD)		1	47	263	197	135	51	4	1	699	382	1,081	80.73	14.69
(%)		(0.1)	(6.7)	(37.6)	(28.2)	(19.3)	(7.3)	(0.6)	(0.1)	(100.0)				
合計	80	796	20,480	111,878	59,687	32,251	15,771	964	75	241,982	68,126	310,108	77.77	14.45
(%)	(0.0)	(0.3)	(8.5)	(46.2)	(24.7)	(13.3)	(6.5)	(0.4)	(0.0)	(100.0)				
記載なし														
(%)														
総計	80	796	20,480	111,878	59,687	32,251	15,771	964	75	241,982	68,126	310,108	77.77	14.45
(%)	(0.0)	(0.3)	(8.5)	(46.2)	(24.7)	(13.3)	(6.5)	(0.4)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査