

表 3831 透析前拡張期血圧(mmHg) 原疾患別（透析患者全体）

原疾患	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	32	265	5,752	33,573	19,719	10,905	5,021	305	15	75,587	21,383	96,970	78.44	14.33
(%)	(0.0)	(0.4)	(7.6)	(44.4)	(26.1)	(14.4)	(6.6)	(0.4)	(0.0)	(100.0)				
慢性腎盂腎炎	1	14	219	1,080	566	310	165	12		2,367	675	3,042	77.60	14.87
(%)	(0.0)	(0.6)	(9.3)	(45.6)	(23.9)	(13.1)	(7.0)	(0.5)		(100.0)				
急速進行性糸球体腎炎		5	164	927	495	262	114	5		1,972	522	2,494	77.65	13.78
(%)		(0.3)	(8.3)	(47.0)	(25.1)	(13.3)	(5.8)	(0.3)		(100.0)				
妊娠腎/妊娠中毒症		9	119	576	343	170	66	7		1,290	324	1,614	77.41	14.37
(%)		(0.7)	(9.2)	(44.7)	(26.6)	(13.2)	(5.1)	(0.5)		(100.0)				
その他の分類不能の腎炎	1	5	85	435	278	176	100	6	1	1,087	286	1,373	79.74	15.32
(%)	(0.1)	(0.5)	(7.8)	(40.0)	(25.6)	(16.2)	(9.2)	(0.6)	(0.1)	(100.0)				
多発性嚢胞腎		12	501	3,848	2,388	1,410	589	18	4	8,770	2,236	11,006	79.55	13.39
(%)		(0.1)	(5.7)	(43.9)	(27.2)	(16.1)	(6.7)	(0.2)	(0.0)	(100.0)				
腎硬化症	7	77	2,308	11,241	5,078	2,443	1,174	81	6	22,415	5,883	28,298	75.99	14.30
(%)	(0.0)	(0.3)	(10.3)	(50.1)	(22.7)	(10.9)	(5.2)	(0.4)	(0.0)	(100.0)				
悪性高血圧		8	116	758	468	330	249	20	3	1,952	621	2,573	81.87	16.01
(%)		(0.4)	(5.9)	(38.8)	(24.0)	(16.9)	(12.8)	(1.0)	(0.2)	(100.0)				
糖尿病性腎症	28	254	7,945	43,993	22,088	11,686	6,004	376	30	92,404	25,677	118,081	77.49	14.41
(%)	(0.0)	(0.3)	(8.6)	(47.6)	(23.9)	(12.6)	(6.5)	(0.4)	(0.0)	(100.0)				
SLE腎炎		7	123	731	444	263	159	7		1,734	502	2,236	79.50	14.73
(%)		(0.4)	(7.1)	(42.2)	(25.6)	(15.2)	(9.2)	(0.4)		(100.0)				
アミロイド腎		1	48	167	81	44	19	1		361	90	451	75.52	14.90
(%)		(0.3)	(13.3)	(46.3)	(22.4)	(12.2)	(5.3)	(0.3)		(100.0)				
痛風腎	2	3	52	420	241	117	50	2		887	226	1,113	77.68	14.00
(%)	(0.2)	(0.3)	(5.9)	(47.4)	(27.2)	(13.2)	(5.6)	(0.2)		(100.0)				
先天性代謝異常による腎不全		1	22	72	56	30	22	4		207	65	272	80.78	16.25
(%)		(0.5)	(10.6)	(34.8)	(27.1)	(14.5)	(10.6)	(1.9)		(100.0)				
腎・尿路結核		3	24	70	29	19	7			152	54	206	73.79	15.81
(%)		(2.0)	(15.8)	(46.1)	(19.1)	(12.5)	(4.6)			(100.0)				
腎・尿路結石		4	43	241	89	64	14			455	126	581	75.57	13.25
(%)		(0.9)	(9.5)	(53.0)	(19.6)	(14.1)	(3.1)			(100.0)				
腎・尿路腫瘍			57	359	194	83	28	1		722	164	886	76.69	12.79
(%)			(7.9)	(49.7)	(26.9)	(11.5)	(3.9)	(0.1)		(100.0)				
閉塞性尿路障害		2	44	226	152	75	44	4		547	176	723	79.05	15.00
(%)		(0.4)	(8.0)	(41.3)	(27.8)	(13.7)	(8.0)	(0.7)		(100.0)				
骨髄腫		2	24	103	56	18	6	1		210	66	276	74.81	13.78
(%)		(1.0)	(11.4)	(49.0)	(26.7)	(8.6)	(2.9)	(0.5)		(100.0)				
腎形成不全		4	39	186	131	87	59	4		510	139	649	81.32	15.95
(%)		(0.8)	(7.6)	(36.5)	(25.7)	(17.1)	(11.6)	(0.8)		(100.0)				
不明	6	85	2,129	9,632	4,897	2,656	1,317	73	10	20,805	6,739	27,544	77.05	14.81
(%)	(0.0)	(0.4)	(10.2)	(46.3)	(23.5)	(12.8)	(6.3)	(0.4)	(0.0)	(100.0)				
再導入	1	6	107	661	463	336	158	13		1,745	418	2,163	80.88	14.88
(%)	(0.1)	(0.3)	(6.1)	(37.9)	(26.5)	(19.3)	(9.1)	(0.7)		(100.0)				
その他	2	29	559	2,579	1,431	767	406	24	6	5,803	1,754	7,557	77.56	15.18
(%)	(0.0)	(0.5)	(9.6)	(44.4)	(24.7)	(13.2)	(7.0)	(0.4)	(0.1)	(100.0)				
合計	80	796	20,480	111,878	59,687	32,251	15,771	964	75	241,982	68,126	310,108	77.77	14.45
(%)	(0.0)	(0.3)	(8.5)	(46.2)	(24.7)	(13.3)	(6.5)	(0.4)	(0.0)	(100.0)				
記載なし														
(%)														
総計	80	796	20,480	111,878	59,687	32,251	15,771	964	75	241,982	68,126	310,108	77.77	14.45
(%)	(0.0)	(0.3)	(8.5)	(46.2)	(24.7)	(13.3)	(6.5)	(0.4)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査