

表 3838 透析前拡張期血圧(mmHg) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	24	195	4,396	25,712	14,818	8,109	3,722	217	13	57,206	11,371	68,577	78.23	14.29
(%)	(0.0)	(0.3)	(7.7)	(44.9)	(25.9)	(14.2)	(6.5)	(0.4)	(0.0)	(100.0)				
慢性腎盂腎炎	1	7	167	818	415	230	117	10		1,765	400	2,165	77.52	14.82
(%)	(0.1)	(0.4)	(9.5)	(46.3)	(23.5)	(13.0)	(6.6)	(0.6)		(100.0)				
急速進行性糸球体腎炎		3	131	727	404	202	92	5		1,564	270	1,834	77.68	13.78
(%)		(0.2)	(8.4)	(46.5)	(25.8)	(12.9)	(5.9)	(0.3)		(100.0)				
妊娠腎/妊娠中毒症		7	75	393	236	120	41	6		878	186	1,064	77.54	14.12
(%)		(0.8)	(8.5)	(44.8)	(26.9)	(13.7)	(4.7)	(0.7)		(100.0)				
その他の分類不能の腎炎	1	3	59	340	195	130	70	5	1	804	145	949	79.49	15.32
(%)	(0.1)	(0.4)	(7.3)	(42.3)	(24.3)	(16.2)	(8.7)	(0.6)	(0.1)	(100.0)				
多発性嚢胞腎		11	393	3,077	1,872	1,109	468	14	3	6,947	1,251	8,198	79.48	13.41
(%)		(0.2)	(5.7)	(44.3)	(26.9)	(16.0)	(6.7)	(0.2)	(0.0)	(100.0)				
腎硬化症	7	64	1,902	9,297	4,082	1,936	909	67	2	18,266	3,194	21,460	75.73	14.19
(%)	(0.0)	(0.4)	(10.4)	(50.9)	(22.3)	(10.6)	(5.0)	(0.4)	(0.0)	(100.0)				
悪性高血圧		7	97	597	359	245	188	15	2	1,510	371	1,881	81.43	16.08
(%)		(0.5)	(6.4)	(39.5)	(23.8)	(16.2)	(12.5)	(1.0)	(0.1)	(100.0)				
糖尿病性腎症	26	203	6,614	36,429	18,069	9,388	4,834	304	24	75,891	14,616	90,507	77.30	14.37
(%)	(0.0)	(0.3)	(8.7)	(48.0)	(23.8)	(12.4)	(6.4)	(0.4)	(0.0)	(100.0)				
SLE腎炎		5	96	553	329	185	120	4		1,292	258	1,550	79.18	14.73
(%)		(0.4)	(7.4)	(42.8)	(25.5)	(14.3)	(9.3)	(0.3)		(100.0)				
アミロイド腎	1	37	134	62	30	13	13	1		278	47	325	74.87	14.79
(%)		(0.4)	(13.3)	(48.2)	(22.3)	(10.8)	(4.7)	(0.4)		(100.0)				
痛風腎	2	3	42	334	191	87	38	2		699	129	828	77.46	14.08
(%)	(0.3)	(0.4)	(6.0)	(47.8)	(27.3)	(12.4)	(5.4)	(0.3)		(100.0)				
先天性代謝異常による腎不全		1	13	54	41	19	16	3		147	30	177	80.82	15.70
(%)		(0.7)	(8.8)	(36.7)	(27.9)	(12.9)	(10.9)	(2.0)		(100.0)				
腎・尿路結核		2	18	56	23	16	7			122	36	158	74.90	15.86
(%)		(1.6)	(14.8)	(45.9)	(18.9)	(13.1)	(5.7)			(100.0)				
腎・尿路結石		3	30	193	74	49	11			360	70	430	75.52	13.15
(%)		(0.8)	(8.3)	(53.6)	(20.6)	(13.6)	(3.1)			(100.0)				
腎・尿路腫瘍			45	309	160	71	24			609	77	686	76.59	12.56
(%)			(7.4)	(50.7)	(26.3)	(11.7)	(3.9)			(100.0)				
閉塞性尿路障害		1	25	181	122	58	37	3		427	90	517	79.73	14.51
(%)		(0.2)	(5.9)	(42.4)	(28.6)	(13.6)	(8.7)	(0.7)		(100.0)				
骨髄腫		2	21	77	42	13	6			161	29	190	74.19	14.10
(%)		(1.2)	(13.0)	(47.8)	(26.1)	(8.1)	(3.7)			(100.0)				
腎形成不全		3	22	125	93	70	41	3		357	62	419	82.05	15.78
(%)		(0.8)	(6.2)	(35.0)	(26.1)	(19.6)	(11.5)	(0.8)		(100.0)				
不明	4	62	1,673	7,718	3,790	2,035	985	54	8	16,329	3,796	20,125	76.81	14.68
(%)	(0.0)	(0.4)	(10.2)	(47.3)	(23.2)	(12.5)	(6.0)	(0.3)	(0.0)	(100.0)				
再導入	1	6	68	457	321	253	110	10		1,226	191	1,417	81.18	14.90
(%)	(0.1)	(0.5)	(5.5)	(37.3)	(26.2)	(20.6)	(9.0)	(0.8)		(100.0)				
その他	1	23	433	2,044	1,142	593	293	18	4	4,551	963	5,514	77.35	14.93
(%)	(0.0)	(0.5)	(9.5)	(44.9)	(25.1)	(13.0)	(6.4)	(0.4)	(0.1)	(100.0)				
合計	67	612	16,357	89,625	46,840	24,948	12,142	741	57	191,389	37,582	228,971	77.56	14.39
(%)	(0.0)	(0.3)	(8.5)	(46.8)	(24.5)	(13.0)	(6.3)	(0.4)	(0.0)	(100.0)				
記載なし														
(%)														
総計	67	612	16,357	89,625	46,840	24,948	12,142	741	57	191,389	37,582	228,971	77.56	14.39
(%)	(0.0)	(0.3)	(8.5)	(46.8)	(24.5)	(13.0)	(6.3)	(0.4)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査