

表 3854 拡張期血圧(mmHg) 体重(kg)別 (腹膜透析患者)

体重	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
20未満		1	10	15	3					29	9	38	61.83	14.12
(%)		(3.4)	(34.5)	(51.7)	(10.3)					(100.0)				
20～			3	10	5	1	2			21	2	23	73.76	17.10
(%)			(14.3)	(47.6)	(23.8)	(4.8)	(9.5)			(100.0)				
30～		3	27	62	31	15	4			142	30	172	72.24	15.40
(%)		(2.1)	(19.0)	(43.7)	(21.8)	(10.6)	(2.8)			(100.0)				
40～		2	76	359	172	87	42	1		739	165	904	76.60	14.09
(%)		(0.3)	(10.3)	(48.6)	(23.3)	(11.8)	(5.7)	(0.1)		(100.0)				
50～		7	100	627	341	185	79	3		1,342	249	1,591	77.91	13.98
(%)		(0.5)	(7.5)	(46.7)	(25.4)	(13.8)	(5.9)	(0.2)		(100.0)				
60～	1	2	92	585	401	201	101	3	1	1,387	229	1,616	79.47	13.88
(%)	(0.1)	(0.1)	(6.6)	(42.2)	(28.9)	(14.5)	(7.3)	(0.2)	(0.1)	(100.0)				
70～			37	257	193	125	61	5		678	133	811	81.70	14.45
(%)			(5.5)	(37.9)	(28.5)	(18.4)	(9.0)	(0.7)		(100.0)				
80～			10	74	70	56	39	2		251	47	298	85.40	14.64
(%)			(4.0)	(29.5)	(27.9)	(22.3)	(15.5)	(0.8)		(100.0)				
90～			6	27	29	20	11	1		94	10	104	84.10	14.56
(%)			(6.4)	(28.7)	(30.9)	(21.3)	(11.7)	(1.1)		(100.0)				
100～			1	4	4	5	1	1		16	4	20	87.50	17.77
(%)			(6.3)	(25.0)	(25.0)	(31.3)	(6.3)	(6.3)		(100.0)				
110～				4	1	2	1			8	2	10	84.38	13.20
(%)				(50.0)	(12.5)	(25.0)	(12.5)			(100.0)				
120～				2	3		1	1		7	2	9	89.29	19.92
(%)				(28.6)	(42.9)		(14.3)	(14.3)		(100.0)				
合計	1	15	362	2,026	1,253	697	342	17	1	4,714	882	5,596	79.00	14.46
(%)	(0.0)	(0.3)	(7.7)	(43.0)	(26.6)	(14.8)	(7.3)	(0.4)	(0.0)	(100.0)				
記載なし			13	39	43	17	5	1		118	3,227	3,345	79.42	13.46
(%)			(11.0)	(33.1)	(36.4)	(14.4)	(4.2)	(0.8)		(100.0)				
総計	1	15	375	2,065	1,296	714	347	18	1	4,832	4,109	8,941	79.01	14.43
(%)	(0.0)	(0.3)	(7.8)	(42.7)	(26.8)	(14.8)	(7.2)	(0.4)	(0.0)	(100.0)				
平均	61.40	48.37	55.93	59.37	62.08	63.68	65.03	77.74	65.60	60.91	59.86	60.74		
標準偏差		14.80	14.88	13.24	13.40	13.44	14.13	27.17		13.83	14.20	13.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査