

表 3861 透析前脈拍(拍/分) 併用の状況別 (透析患者全体)

併用の状況	40未満	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	180	2,354	19,300	59,869	71,918	45,728	20,087	6,179	1,474	634	227,723	73,071	300,794	74.63	12.82
(%)	(0.1)	(1.0)	(8.5)	(26.3)	(31.6)	(20.1)	(8.8)	(2.7)	(0.6)	(0.3)	(100.0)				
PD単独	5	11	112	568	907	755	355	154	40	18	2,925	4,263	7,188	78.78	13.28
(%)	(0.2)	(0.4)	(3.8)	(19.4)	(31.0)	(25.8)	(12.1)	(5.3)	(1.4)	(0.6)	(100.0)				
HD(F)単独.PDカテあり		2	18	51	55	36	15	8	1	1	187	82	269	74.53	13.75
(%)		(1.1)	(9.6)	(27.3)	(29.4)	(19.3)	(8.0)	(4.3)	(0.5)	(0.5)	(100.0)				
PD.HD併用週1		12	67	255	321	245	110	47	11	2	1,070	474	1,544	76.41	13.13
HD(F)															
(%)		(1.1)	(6.3)	(23.8)	(30.0)	(22.9)	(10.3)	(4.4)	(1.0)	(0.2)	(100.0)				
PD.HD併用週2		2	7	18	40	24	9	4	1		105	72	177	76.33	12.61
HD(F)															
(%)		(1.9)	(6.7)	(17.1)	(38.1)	(22.9)	(8.6)	(3.8)	(1.0)		(100.0)				
PD.HD併用週3			2	4	12	6	4		1		29	11	40	78.38	12.41
HD(F)															
(%)			(6.9)	(13.8)	(41.4)	(20.7)	(13.8)		(3.4)		(100.0)				
PD.HD併用週4				1							1		1	66.00	
HD(F)															
(%)				(100.0)							(100.0)				
PD.HD併用その他			3	14	17	11	10	1			56	17	73	77.25	11.90
(%)			(5.4)	(25.0)	(30.4)	(19.6)	(17.9)	(1.8)			(100.0)				
合計	185	2,381	19,509	60,780	73,270	46,805	20,590	6,393	1,528	655	232,096	77,990	310,086	74.69	12.84
(%)	(0.1)	(1.0)	(8.4)	(26.2)	(31.6)	(20.2)	(8.9)	(2.8)	(0.7)	(0.3)	(100.0)				
不明			3	1	2	9					15	7	22	75.93	12.69
(%)			(20.0)	(6.7)	(13.3)	(60.0)					(100.0)				
記載なし															
(%)															
総計	185	2,381	19,512	60,781	73,272	46,814	20,590	6,393	1,528	655	232,111	77,997	310,108	74.69	12.84
(%)	(0.1)	(1.0)	(8.4)	(26.2)	(31.6)	(20.2)	(8.9)	(2.8)	(0.7)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査