

表 3917 PET施行の有無 原疾患別（腹膜透析単独患者）

原疾患	なし	PET施行	Fast PETのみ	合計	不明	記載なし	総計
慢性糸球体腎炎	455	658	351	1,464	28	837	2,329
(%)	(31.1)	(44.9)	(24.0)	(100.0)			
慢性腎盂腎炎	14	24	11	49	1	29	79
(%)	(28.6)	(49.0)	(22.4)	(100.0)			
急速進行性糸球体腎炎	10	20	7	37		27	64
(%)	(27.0)	(54.1)	(18.9)	(100.0)			
妊娠腎/妊娠中毒症	2	12	3	17		5	22
(%)	(11.8)	(70.6)	(17.6)	(100.0)			
その他の分類不能の腎炎	13	19	9	41	2	23	66
(%)	(31.7)	(46.3)	(22.0)	(100.0)			
多発性嚢胞腎	52	56	25	133		73	206
(%)	(39.1)	(42.1)	(18.8)	(100.0)			
腎硬化症	246	221	127	594	18	328	940
(%)	(41.4)	(37.2)	(21.4)	(100.0)			
悪性高血圧	17	23	11	51		34	85
(%)	(33.3)	(45.1)	(21.6)	(100.0)			
糖尿病性腎症	460	557	293	1,310	27	898	2,235
(%)	(35.1)	(42.5)	(22.4)	(100.0)			
SLE腎炎	13	13	5	31	1	22	54
(%)	(41.9)	(41.9)	(16.1)	(100.0)			
アミロイド腎	2	1	1	4		3	7
(%)	(50.0)	(25.0)	(25.0)	(100.0)			
痛風腎	7	7	5	19		7	26
(%)	(36.8)	(36.8)	(26.3)	(100.0)			
先天性代謝異常による腎不全	4	2	4	10	1	6	17
(%)	(40.0)	(20.0)	(40.0)	(100.0)			
腎・尿路結核		1		1			1
(%)		(100.0)		(100.0)			
腎・尿路結石	2	5	1	8		6	14
(%)	(25.0)	(62.5)	(12.5)	(100.0)			
腎・尿路腫瘍	3	4	2	9		2	11
(%)	(33.3)	(44.4)	(22.2)	(100.0)			
閉塞性尿路障害	3	7	2	12		7	19
(%)	(25.0)	(58.3)	(16.7)	(100.0)			
骨髄腫	2	1		3		4	7
(%)	(66.7)	(33.3)		(100.0)			
腎形成不全	18	17	9	44		28	72
(%)	(40.9)	(38.6)	(20.5)	(100.0)			
不明	143	168	93	404	10	275	689
(%)	(35.4)	(41.6)	(23.0)	(100.0)			
再導入	5	17	4	26	2	8	36
(%)	(19.2)	(65.4)	(15.4)	(100.0)			
その他	42	52	29	123		86	209
(%)	(34.1)	(42.3)	(23.6)	(100.0)			
合計	1,513	1,885	992	4,390	90	2,708	7,188
(%)	(34.5)	(42.9)	(22.6)	(100.0)			
記載なし							
(%)							
総計	1,513	1,885	992	4,390	90	2,708	7,188
(%)	(34.5)	(42.9)	(22.6)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査