

表 3951 一日透析液使用量(L/日) 原疾患別(腹膜透析患者)

原疾患	2未満	2～	4～	6～	8～	10～	12～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	30	99	334	656	557	165	80	1,921	1,138	3,059	7.08	2.71
(%)	(1.6)	(5.2)	(17.4)	(34.1)	(29.0)	(8.6)	(4.2)	(100.0)				
慢性腎盂腎炎	4	7	14	21	9	7	2	64	29	93	6.16	2.81
(%)	(6.3)	(10.9)	(21.9)	(32.8)	(14.1)	(10.9)	(3.1)	(100.0)				
急速進行性糸球体腎炎	1	2	7	15	13	1	4	43	31	74	7.30	2.80
(%)	(2.3)	(4.7)	(16.3)	(34.9)	(30.2)	(2.3)	(9.3)	(100.0)				
妊娠腎/妊娠中毒症	2		6	11	3	2		24	8	32	6.13	2.42
(%)	(8.3)		(25.0)	(45.8)	(12.5)	(8.3)		(100.0)				
その他の分類不能の腎炎		4	13	12	19	4	3	55	30	85	7.21	2.54
(%)		(7.3)	(23.6)	(21.8)	(34.5)	(7.3)	(5.5)	(100.0)				
多発性嚢胞腎	6	17	40	38	37	10	5	153	90	243	6.39	2.89
(%)	(3.9)	(11.1)	(26.1)	(24.8)	(24.2)	(6.5)	(3.3)	(100.0)				
腎硬化症	12	79	160	231	142	37	12	673	410	1,083	6.25	2.46
(%)	(1.8)	(11.7)	(23.8)	(34.3)	(21.1)	(5.5)	(1.8)	(100.0)				
悪性高血圧	2	8	12	22	15	4	1	64	43	107	6.37	2.45
(%)	(3.1)	(12.5)	(18.8)	(34.4)	(23.4)	(6.3)	(1.6)	(100.0)				
糖尿病性腎症	34	102	328	532	414	124	57	1,591	1,175	2,766	6.79	2.65
(%)	(2.1)	(6.4)	(20.6)	(33.4)	(26.0)	(7.8)	(3.6)	(100.0)				
SLE腎炎		4	5	14	10	2	2	37	25	62	7.04	2.95
(%)		(10.8)	(13.5)	(37.8)	(27.0)	(5.4)	(5.4)	(100.0)				
アミロイド腎	1	1	1	1	1	1		5	4	9	6.72	2.98
(%)		(20.0)	(20.0)	(20.0)	(20.0)	(20.0)		(100.0)				
痛風腎		1	4	4	11	2	1	23	10	33	7.63	2.33
(%)		(4.3)	(17.4)	(17.4)	(47.8)	(8.7)	(4.3)	(100.0)				
先天性代謝異常による腎不全	1	1	3	4	3			12	6	18	5.55	2.21
(%)	(8.3)	(8.3)	(25.0)	(33.3)	(25.0)			(100.0)				
腎・尿路結核				1				1		1	6.00	
(%)				(100.0)				(100.0)				
腎・尿路結石			1	2	6			9	7	16	7.56	1.45
(%)			(11.1)	(22.2)	(66.7)			(100.0)				
腎・尿路腫瘍			2	3	3	1		9	5	14	7.10	1.88
(%)			(22.2)	(33.3)	(33.3)	(11.1)		(100.0)				
閉塞性尿路障害		3		3	4	1	1	12	10	22	6.91	2.61
(%)		(25.0)		(25.0)	(33.3)	(8.3)	(8.3)	(100.0)				
骨髄腫			1	1	1			3	5	8	6.50	2.29
(%)			(33.3)	(33.3)	(33.3)			(100.0)				
腎形成不全	5	7	10	4	12	3	6	47	29	76	6.86	4.52
(%)	(10.6)	(14.9)	(21.3)	(8.5)	(25.5)	(6.4)	(12.8)	(100.0)				
不明	13	37	95	162	110	34	18	469	371	840	6.60	2.73
(%)	(2.8)	(7.9)	(20.3)	(34.5)	(23.5)	(7.2)	(3.8)	(100.0)				
再導入		2	4	15	10	4		35	16	51	6.96	1.95
(%)		(5.7)	(11.4)	(42.9)	(28.6)	(11.4)		(100.0)				
その他	9	19	32	43	26	8	9	146	103	249	6.26	3.43
(%)	(6.2)	(13.0)	(21.9)	(29.5)	(17.8)	(5.5)	(6.2)	(100.0)				
合計	119	393	1,072	1,795	1,406	410	201	5,396	3,545	8,941	6.78	2.71
(%)	(2.2)	(7.3)	(19.9)	(33.3)	(26.1)	(7.6)	(3.7)	(100.0)				
記載なし												
(%)												
総計	119	393	1,072	1,795	1,406	410	201	5,396	3,545	8,941	6.78	2.71
(%)	(2.2)	(7.3)	(19.9)	(33.3)	(26.1)	(7.6)	(3.7)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査