

表 3984 残存腎機能(一日尿量)(mL/日) 原疾患別 (PD・HD(F)等併用患者)

原疾患	100未満	100～	400～	800～	1200～	1600～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	215	73	34	16	1	2	341	386	727	153.87	287.87
(%)	(63.0)	(21.4)	(10.0)	(4.7)	(0.3)	(0.6)	(100.0)				
慢性腎盂腎炎	7	4					11	3	14	66.36	105.10
(%)	(63.6)	(36.4)					(100.0)				
急速進行性糸球体腎炎	4	1					5	5	10	40.00	89.44
(%)	(80.0)	(20.0)					(100.0)				
妊娠腎/妊娠中毒症	4	1					5	5	10	54.00	87.06
(%)	(80.0)	(20.0)					(100.0)				
その他の分類不能の腎炎	7	1	1				9	10	19	67.22	165.59
(%)	(77.8)	(11.1)	(11.1)				(100.0)				
多発性嚢胞腎	9	6	2	3		1	21	16	37	350.71	504.19
(%)	(42.9)	(28.6)	(9.5)	(14.3)		(4.8)	(100.0)				
腎硬化症	32	18	3	5	1	1	60	83	143	207.00	363.57
(%)	(53.3)	(30.0)	(5.0)	(8.3)	(1.7)	(1.7)	(100.0)				
悪性高血圧	9	1	2	1			13	9	22	192.31	340.86
(%)	(69.2)	(7.7)	(15.4)	(7.7)			(100.0)				
糖尿病性腎症	110	64	27	14	5	2	222	308	530	231.38	356.45
(%)	(49.5)	(28.8)	(12.2)	(6.3)	(2.3)	(0.9)	(100.0)				
SLE腎炎	3				1		4	4	8	300.00	600.00
(%)	(75.0)				(25.0)		(100.0)				
アミロイド腎								2	2		
(%)											
痛風腎	3	2					5	2	7	60.00	89.44
(%)	(60.0)	(40.0)					(100.0)				
先天性代謝異常による腎不全	1						1		1		
(%)	(100.0)						(100.0)				
腎・尿路結核											
(%)											
腎・尿路結石		1					1	1	2	150.00	
(%)		(100.0)					(100.0)				
腎・尿路腫瘍	1						1	2	3		
(%)	(100.0)						(100.0)				
閉塞性尿路障害	1						1	2	3		
(%)	(100.0)						(100.0)				
骨髄腫								1	1		
(%)											
腎形成不全	1	1					2	2	4	175.00	219.20
(%)	(50.0)	(50.0)					(100.0)				
不明	42	10	4	5	1		62	88	150	176.60	336.81
(%)	(67.7)	(16.1)	(6.5)	(8.1)	(1.6)		(100.0)				
再導入	8						8	7	15		
(%)	(100.0)						(100.0)				
その他	7	6	2				15	25	40	155.33	177.88
(%)	(46.7)	(40.0)	(13.3)				(100.0)				
合計	464	189	75	44	9	6	787	961	1,748	181.96	322.70
(%)	(59.0)	(24.0)	(9.5)	(5.6)	(1.1)	(0.8)	(100.0)				
記載なし											
(%)											
総計	464	189	75	44	9	6	787	961	1,748	181.96	322.70
(%)	(59.0)	(24.0)	(9.5)	(5.6)	(1.1)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査