

表 4011 APD使用の有無 原疾患別（腹膜透析患者）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,075	878	1,953	7	1,099	3,059
(%)	(55.0)	(45.0)	(100.0)			
慢性腎盂腎炎	30	28	58	1	34	93
(%)	(51.7)	(48.3)	(100.0)			
急速進行性糸球体腎炎	14	29	43		31	74
(%)	(32.6)	(67.4)	(100.0)			
妊娠腎/妊娠中毒症	14	10	24		8	32
(%)	(58.3)	(41.7)	(100.0)			
その他の分類不能の腎炎	19	33	52	1	32	85
(%)	(36.5)	(63.5)	(100.0)			
多発性嚢胞腎	75	78	153	2	88	243
(%)	(49.0)	(51.0)	(100.0)			
腎硬化症	375	312	687	5	391	1,083
(%)	(54.6)	(45.4)	(100.0)			
悪性高血圧	29	37	66		41	107
(%)	(43.9)	(56.1)	(100.0)			
糖尿病性腎症	887	721	1,608	12	1,146	2,766
(%)	(55.2)	(44.8)	(100.0)			
SLE腎炎	18	19	37		25	62
(%)	(48.6)	(51.4)	(100.0)			
アミロイド腎	3	1	4		5	9
(%)	(75.0)	(25.0)	(100.0)			
痛風腎	12	12	24		9	33
(%)	(50.0)	(50.0)	(100.0)			
先天性代謝異常による腎不全	7	6	13		5	18
(%)	(53.8)	(46.2)	(100.0)			
腎・尿路結核	1		1			1
(%)	(100.0)		(100.0)			
腎・尿路結石	6	3	9		7	16
(%)	(66.7)	(33.3)	(100.0)			
腎・尿路腫瘍	3	6	9		5	14
(%)	(33.3)	(66.7)	(100.0)			
閉塞性尿路障害	7	6	13		9	22
(%)	(53.8)	(46.2)	(100.0)			
骨髄腫	2	1	3		5	8
(%)	(66.7)	(33.3)	(100.0)			
腎形成不全	13	35	48		28	76
(%)	(27.1)	(72.9)	(100.0)			
不明	272	209	481	7	352	840
(%)	(56.5)	(43.5)	(100.0)			
再導入	17	18	35		16	51
(%)	(48.6)	(51.4)	(100.0)			
その他	77	68	145		104	249
(%)	(53.1)	(46.9)	(100.0)			
合計	2,956	2,510	5,466	35	3,440	8,941
(%)	(54.1)	(45.9)	(100.0)			
記載なし						
(%)						
総計	2,956	2,510	5,466	35	3,440	8,941
(%)	(54.1)	(45.9)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査