

表 4017 APD使用の有無 原疾患別（腹膜透析単独患者）

| 原疾患           | なし      | あり     | 合計      | 不明 | 記載なし  | 総計    |
|---------------|---------|--------|---------|----|-------|-------|
| 慢性糸球体腎炎       | 840     | 654    | 1,494   | 7  | 828   | 2,329 |
| (%)           | (56.2)  | (43.8) | (100.0) |    |       |       |
| 慢性腎盂腎炎        | 24      | 23     | 47      | 1  | 31    | 79    |
| (%)           | (51.1)  | (48.9) | (100.0) |    |       |       |
| 急速進行性糸球体腎炎    | 10      | 27     | 37      |    | 27    | 64    |
| (%)           | (27.0)  | (73.0) | (100.0) |    |       |       |
| 妊娠腎/妊娠中毒症     | 9       | 8      | 17      |    | 5     | 22    |
| (%)           | (52.9)  | (47.1) | (100.0) |    |       |       |
| その他の分類不能の腎炎   | 15      | 26     | 41      | 1  | 24    | 66    |
| (%)           | (36.6)  | (63.4) | (100.0) |    |       |       |
| 多発性嚢胞腎        | 64      | 66     | 130     | 1  | 75    | 206   |
| (%)           | (49.2)  | (50.8) | (100.0) |    |       |       |
| 腎硬化症          | 336     | 272    | 608     | 4  | 328   | 940   |
| (%)           | (55.3)  | (44.7) | (100.0) |    |       |       |
| 悪性高血圧         | 23      | 28     | 51      |    | 34    | 85    |
| (%)           | (45.1)  | (54.9) | (100.0) |    |       |       |
| 糖尿病性腎症        | 746     | 578    | 1,324   | 10 | 901   | 2,235 |
| (%)           | (56.3)  | (43.7) | (100.0) |    |       |       |
| SLE腎炎         | 16      | 16     | 32      |    | 22    | 54    |
| (%)           | (50.0)  | (50.0) | (100.0) |    |       |       |
| アミロイド腎        | 3       | 1      | 4       |    | 3     | 7     |
| (%)           | (75.0)  | (25.0) | (100.0) |    |       |       |
| 痛風腎           | 7       | 11     | 18      |    | 8     | 26    |
| (%)           | (38.9)  | (61.1) | (100.0) |    |       |       |
| 先天性代謝異常による腎不全 | 7       | 5      | 12      |    | 5     | 17    |
| (%)           | (58.3)  | (41.7) | (100.0) |    |       |       |
| 腎・尿路結核        | 1       |        | 1       |    |       | 1     |
| (%)           | (100.0) |        | (100.0) |    |       |       |
| 腎・尿路結石        | 6       | 2      | 8       |    | 6     | 14    |
| (%)           | (75.0)  | (25.0) | (100.0) |    |       |       |
| 腎・尿路腫瘍        | 3       | 4      | 7       |    | 4     | 11    |
| (%)           | (42.9)  | (57.1) | (100.0) |    |       |       |
| 閉塞性尿路障害       | 7       | 5      | 12      |    | 7     | 19    |
| (%)           | (58.3)  | (41.7) | (100.0) |    |       |       |
| 骨髄腫           | 2       | 1      | 3       |    | 4     | 7     |
| (%)           | (66.7)  | (33.3) | (100.0) |    |       |       |
| 腎形成不全         | 12      | 33     | 45      |    | 27    | 72    |
| (%)           | (26.7)  | (73.3) | (100.0) |    |       |       |
| 不明            | 225     | 180    | 405     | 6  | 278   | 689   |
| (%)           | (55.6)  | (44.4) | (100.0) |    |       |       |
| 再導入           | 14      | 12     | 26      |    | 10    | 36    |
| (%)           | (53.8)  | (46.2) | (100.0) |    |       |       |
| その他           | 68      | 56     | 124     |    | 85    | 209   |
| (%)           | (54.8)  | (45.2) | (100.0) |    |       |       |
| 合計            | 2,438   | 2,008  | 4,446   | 30 | 2,712 | 7,188 |
| (%)           | (54.8)  | (45.2) | (100.0) |    |       |       |
| 記載なし          |         |        |         |    |       |       |
| (%)           |         |        |         |    |       |       |
| 総計            | 2,438   | 2,008  | 4,446   | 30 | 2,712 | 7,188 |
| (%)           | (54.8)  | (45.2) | (100.0) |    |       |       |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査