

表 4029 1日のPD実施時間(時間/日) 年齢(歳)別(腹膜透析患者)

| 年齢    | 1～    | 5～     | 9～     | 13～    | 18～   | 24     | 合計      | 記載なし  | 総計    | 平均    | 標準偏差 |
|-------|-------|--------|--------|--------|-------|--------|---------|-------|-------|-------|------|
| 15歳未満 | 2     | 14     | 26     | 2      |       | 6      | 50      | 47    | 97    | 11.22 | 5.31 |
| (%)   | (4.0) | (28.0) | (52.0) | (4.0)  |       | (12.0) | (100.0) |       |       |       |      |
| 15歳～  | 2     | 15     | 24     | 1      | 1     | 27     | 70      | 41    | 111   | 14.80 | 7.67 |
| (%)   | (2.9) | (21.4) | (34.3) | (1.4)  | (1.4) | (38.6) | (100.0) |       |       |       |      |
| 30歳～  | 3     | 68     | 46     | 17     | 16    | 256    | 406     | 295   | 701   | 19.01 | 6.98 |
| (%)   | (0.7) | (16.7) | (11.3) | (4.2)  | (3.9) | (63.1) | (100.0) |       |       |       |      |
| 45歳～  | 19    | 219    | 147    | 85     | 78    | 798    | 1,346   | 956   | 2,302 | 18.70 | 7.03 |
| (%)   | (1.4) | (16.3) | (10.9) | (6.3)  | (5.8) | (59.3) | (100.0) |       |       |       |      |
| 60歳～  | 41    | 347    | 287    | 196    | 90    | 1,420  | 2,381   | 1,653 | 4,034 | 18.70 | 6.97 |
| (%)   | (1.7) | (14.6) | (12.1) | (8.2)  | (3.8) | (59.6) | (100.0) |       |       |       |      |
| 75歳～  | 28    | 153    | 166    | 98     | 28    | 461    | 934     | 657   | 1,591 | 17.16 | 7.32 |
| (%)   | (3.0) | (16.4) | (17.8) | (10.5) | (3.0) | (49.4) | (100.0) |       |       |       |      |
| 90歳～  | 3     | 16     | 5      | 4      |       | 36     | 64      | 41    | 105   | 17.25 | 7.98 |
| (%)   | (4.7) | (25.0) | (7.8)  | (6.3)  |       | (56.3) | (100.0) |       |       |       |      |
| 合計    | 98    | 832    | 701    | 403    | 213   | 3,004  | 5,251   | 3,690 | 8,941 | 18.31 | 7.13 |
| (%)   | (1.9) | (15.8) | (13.3) | (7.7)  | (4.1) | (57.2) | (100.0) |       |       |       |      |
| 不明    |       |        |        |        |       |        |         |       |       |       |      |
| (%)   |       |        |        |        |       |        |         |       |       |       |      |
| 記載なし  |       |        |        |        |       |        |         |       |       |       |      |
| (%)   |       |        |        |        |       |        |         |       |       |       |      |
| 総計    | 98    | 832    | 701    | 403    | 213   | 3,004  | 5,251   | 3,690 | 8,941 | 18.31 | 7.13 |
| (%)   | (1.9) | (15.8) | (13.3) | (7.7)  | (4.1) | (57.2) | (100.0) |       |       |       |      |
| 平均    | 65.60 | 61.91  | 61.83  | 66.09  | 60.93 | 62.37  | 62.51   | 62.36 | 62.45 |       |      |
| 標準偏差  | 16.67 | 15.82  | 18.47  | 12.73  | 11.87 | 13.04  | 14.39   | 14.74 | 14.53 |       |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査