

表 4044 1日のPD実施時間(時間/日) 原疾患別 (PD・HD(F)等併用患者)

原疾患	1～	5～	9～	13～	18～	24	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	11	44	29	23	26	299	432	295	727	20.07	6.56
(%)	(2.5)	(10.2)	(6.7)	(5.3)	(6.0)	(69.2)	(100.0)				
慢性腎盂腎炎				2	1	8	11	3	14	21.91	4.04
(%)				(18.2)	(9.1)	(72.7)	(100.0)				
急速進行性糸球体腎炎					1	5	6	4	10	23.00	2.45
(%)					(16.7)	(83.3)	(100.0)				
妊娠腎/妊娠中毒症	1					6	7	3	10	21.00	7.94
(%)	(14.3)					(85.7)	(100.0)				
その他の分類不能の腎炎		1		2	1	7	11	8	19	20.64	5.35
(%)		(9.1)		(18.2)	(9.1)	(63.6)	(100.0)				
多発性嚢胞腎		3	1	5	1	14	24	13	37	19.21	6.48
(%)		(12.5)	(4.2)	(20.8)	(4.2)	(58.3)	(100.0)				
腎硬化症	2	7	12	6	4	46	77	66	143	18.88	6.95
(%)	(2.6)	(9.1)	(15.6)	(7.8)	(5.2)	(59.7)	(100.0)				
悪性高血圧		4		1	1	9	15	7	22	18.53	7.62
(%)		(26.7)		(6.7)	(6.7)	(60.0)	(100.0)				
糖尿病性腎症	2	28	34	19	24	154	261	269	530	19.21	6.58
(%)	(0.8)	(10.7)	(13.0)	(7.3)	(9.2)	(59.0)	(100.0)				
SLE腎炎		1				4	5	3	8	20.80	7.16
(%)		(20.0)				(80.0)	(100.0)				
アミロイド腎								2	2		
(%)											
痛風腎	1		1			4	6	1	7	18.67	8.64
(%)	(16.7)		(16.7)			(66.7)	(100.0)				
先天性代謝異常による腎不全						1	1		1	24.00	
(%)						(100.0)	(100.0)				
腎・尿路結核											
(%)											
腎・尿路結石				1			1	1	2	14.00	
(%)				(100.0)			(100.0)				
腎・尿路腫瘍		1				1	2	1	3	16.00	11.31
(%)		(50.0)				(50.0)	(100.0)				
閉塞性尿路障害						1	1	2	3	24.00	
(%)						(100.0)	(100.0)				
骨髄腫								1	1		
(%)											
腎形成不全		1				1	2	2	4	16.00	11.31
(%)		(50.0)				(50.0)	(100.0)				
不明	2	6	6	3	5	51	73	77	150	20.22	6.52
(%)	(2.7)	(8.2)	(8.2)	(4.1)	(6.8)	(69.9)	(100.0)				
再導入			3		1	5	9	6	15	18.78	6.96
(%)			(33.3)		(11.1)	(55.6)	(100.0)				
その他		2	2	1		16	21	19	40	20.67	6.25
(%)		(9.5)	(9.5)	(4.8)		(76.2)	(100.0)				
合計	19	98	88	63	65	632	965	783	1,748	19.74	6.58
(%)	(2.0)	(10.2)	(9.1)	(6.5)	(6.7)	(65.5)	(100.0)				
記載なし											
(%)											
総計	19	98	88	63	65	632	965	783	1,748	19.74	6.58
(%)	(2.0)	(10.2)	(9.1)	(6.5)	(6.7)	(65.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査