

表 4051 PD透析液交換方法 原疾患別（腹膜透析患者）

原疾患	完全手動式	バッグ交換デバイス(紫 外線)	バッグ交換デバイス(熱 式無菌接合装置)	バッグ交換デバイス (左記以外、半手動も 含む)	合計	不明	記載なし	総計
慢性糸球体腎炎	700	978	247	25	1,950	30	1,079	3,059
(%)	(35.9)	(50.2)	(12.7)	(1.3)	(100.0)			
慢性腎盂腎炎	25	24	9		58	2	33	93
(%)	(43.1)	(41.4)	(15.5)		(100.0)			
急速進行性糸球 体腎炎	10	22	10		42	1	31	74
(%)	(23.8)	(52.4)	(23.8)		(100.0)			
妊娠腎/妊娠中毒 症	11	8	5		24		8	32
(%)	(45.8)	(33.3)	(20.8)		(100.0)			
その他の分類不 能の腎炎	20	30	4		54	2	29	85
(%)	(37.0)	(55.6)	(7.4)		(100.0)			
多発性嚢胞腎	39	83	27	5	154	3	86	243
(%)	(25.3)	(53.9)	(17.5)	(3.2)	(100.0)			
腎硬化症	212	349	112	8	681	14	388	1,083
(%)	(31.1)	(51.2)	(16.4)	(1.2)	(100.0)			
悪性高血圧	20	40	6		66		41	107
(%)	(30.3)	(60.6)	(9.1)		(100.0)			
糖尿病性腎症	444	929	182	30	1,585	39	1,142	2,766
(%)	(28.0)	(58.6)	(11.5)	(1.9)	(100.0)			
SLE腎炎	13	16	5	1	35	2	25	62
(%)	(37.1)	(45.7)	(14.3)	(2.9)	(100.0)			
アミロイド腎	2	2	1		5		4	9
(%)	(40.0)	(40.0)	(20.0)		(100.0)			
痛風腎	10	8	4	1	23	1	9	33
(%)	(43.5)	(34.8)	(17.4)	(4.3)	(100.0)			
先天性代謝異常 による腎不全	2	9	1		12		6	18
(%)	(16.7)	(75.0)	(8.3)		(100.0)			
腎・尿路結核	1				1			1
(%)	(100.0)				(100.0)			
腎・尿路結石	5	3	1		9		7	16
(%)	(55.6)	(33.3)	(11.1)		(100.0)			
腎・尿路腫瘍	5	3	1	1	10	1	3	14
(%)	(50.0)	(30.0)	(10.0)	(10.0)	(100.0)			
閉塞性尿路障害	4	5	4		13	1	8	22
(%)	(30.8)	(38.5)	(30.8)		(100.0)			
骨髄腫		3			3		5	8
(%)		(100.0)			(100.0)			
腎形成不全	17	23	7	1	48		28	76
(%)	(35.4)	(47.9)	(14.6)	(2.1)	(100.0)			
不明	165	250	59	7	481	10	349	840
(%)	(34.3)	(52.0)	(12.3)	(1.5)	(100.0)			
再導入	11	18	7		36		15	51
(%)	(30.6)	(50.0)	(19.4)		(100.0)			
その他	44	72	25	1	142	1	106	249
(%)	(31.0)	(50.7)	(17.6)	(0.7)	(100.0)			
合計	1,760	2,875	717	80	5,432	107	3,402	8,941
(%)	(32.4)	(52.9)	(13.2)	(1.5)	(100.0)			
記載なし								
(%)								
総計	1,760	2,875	717	80	5,432	107	3,402	8,941
(%)	(32.4)	(52.9)	(13.2)	(1.5)	(100.0)			

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査