

表 4077 2014年中の腹膜炎発症率(回/1患者・年) 原疾患別(腹膜透析単独患者)

原疾患	0	1.0～	2.0～	3.0～	4.0～	5.0～	合計	不明 記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,290	125	23	11	2	6	1,457	872	2,329	0.19	0.78
(%)	(88.5)	(8.6)	(1.6)	(0.8)	(0.1)	(0.4)	(100.0)				
慢性腎盂腎炎	40	6	2				48	31	79	0.21	0.50
(%)	(83.3)	(12.5)	(4.2)				(100.0)				
急速進行性糸球 体腎炎	30	4	1				35	29	64	0.18	0.50
(%)	(85.7)	(11.4)	(2.9)				(100.0)				
妊娠腎/妊娠中毒 症	14	3					17	5	22	0.18	0.39
(%)	(82.4)	(17.6)					(100.0)				
その他の分類不能 の腎炎	38	2				2	42	24	66	0.35	1.39
(%)	(90.5)	(4.8)				(4.8)	(100.0)				
多発性嚢胞腎	107	12	4	3			126	80	206	0.25	0.66
(%)	(84.9)	(9.5)	(3.2)	(2.4)			(100.0)				
腎硬化症	498	68	12	1	1	4	584	356	940	0.22	0.76
(%)	(85.3)	(11.6)	(2.1)	(0.2)	(0.2)	(0.7)	(100.0)				
悪性高血圧	45	6					51	34	85	0.13	0.36
(%)	(88.2)	(11.8)					(100.0)				
糖尿病性腎症	1,113	106	22	5	9	9	1,264	971	2,235	0.22	0.87
(%)	(88.1)	(8.4)	(1.7)	(0.4)	(0.7)	(0.7)	(100.0)				
SLE腎炎	28	1	3				32	22	54	0.23	0.65
(%)	(87.5)	(3.1)	(9.4)				(100.0)				
アミロイド腎	4						4	3	7		
(%)	(100.0)						(100.0)				
痛風腎	15	2					17	9	26	0.12	0.33
(%)	(88.2)	(11.8)					(100.0)				
先天性代謝異常に よる腎不全	10	1					11	6	17	0.09	0.30
(%)	(90.9)	(9.1)					(100.0)				
腎・尿路結核	1						1		1		
(%)	(100.0)						(100.0)				
腎・尿路結石	7			1			8	6	14	0.38	1.06
(%)	(87.5)			(12.5)			(100.0)				
腎・尿路腫瘍	4	1	1	1			7	4	11	0.87	1.22
(%)	(57.1)	(14.3)	(14.3)	(14.3)			(100.0)				
閉塞性尿路障害	7	3		1			11	8	19	0.56	0.95
(%)	(63.6)	(27.3)		(9.1)			(100.0)				
骨髄腫	2		1				3	4	7	0.67	1.15
(%)	(66.7)		(33.3)				(100.0)				
腎形成不全	40	4					44	28	72	0.09	0.29
(%)	(90.9)	(9.1)					(100.0)				
不明	347	38	12	1		1	399	290	689	0.18	0.55
(%)	(87.0)	(9.5)	(3.0)	(0.3)		(0.3)	(100.0)				
再導入	20	3			1		24	12	36	0.29	0.86
(%)	(83.3)	(12.5)			(4.2)		(100.0)				
その他	98	15	1	1		1	116	93	209	0.28	1.20
(%)	(84.5)	(12.9)	(0.9)	(0.9)		(0.9)	(100.0)				
合計	3,758	400	82	25	13	23	4,301	2,887	7,188	0.21	0.79
(%)	(87.4)	(9.3)	(1.9)	(0.6)	(0.3)	(0.5)	(100.0)				
記載なし (%)											
総計	3,758	400	82	25	13	23	4,301	2,887	7,188	0.21	0.79
(%)	(87.4)	(9.3)	(1.9)	(0.6)	(0.3)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

\*個々の患者の腹膜炎発症率(回/1患者・年)は、1.0未満の数値が0となるため、「0」と表示しています。

(C)Japanese Society for Dialysis Therapy

2014年調査