

表 4084 2014年中の腹膜炎発症率(回/1患者・年) 原疾患別(PD・HD(F)等併用患者)

原疾患	0	1.0~	2.0~	3.0~	4.0~	5.0~	合計	不明 記載なし	総計	平均	標準偏差
慢性糸球体腎炎	396	40	5	3	1	2	447	280	727	0.17	0.66
(%)	(88.6)	(8.9)	(1.1)	(0.7)	(0.2)	(0.4)	(100.0)				
慢性腎孟腎炎	10	1					11	3	14	0.09	0.30
(%)	(90.9)	(9.1)					(100.0)				
急速進行性糸球体腎炎	6						6	4	10		
(%)	(100.0)						(100.0)				
妊娠腎/妊娠中毒症	5	1	1				7	3	10	0.43	0.79
(%)	(71.4)	(14.3)	(14.3)				(100.0)				
その他の分類不能の腎炎	11						11	8	19		
(%)	(100.0)						(100.0)				
多発性囊胞腎	19	3	1				23	14	37	0.23	0.58
(%)	(82.6)	(13.0)	(4.3)				(100.0)				
腎硬化症	67	7	2				76	67	143	0.14	0.42
(%)	(88.2)	(9.2)	(2.6)				(100.0)				
悪性高血圧	14						14	8	22		
(%)	(100.0)						(100.0)				
糖尿病性腎症	249	24	3	1	1		278	252	530	0.14	0.46
(%)	(89.6)	(8.6)	(1.1)	(0.4)	(0.4)		(100.0)				
SLE腎炎	5						5	3	8		
(%)	(100.0)						(100.0)				
アミロイド腎	1						1	1	2		
(%)	(100.0)						(100.0)				
痛風腎	6						6	1	7		
(%)	(100.0)						(100.0)				
先天性代謝異常に による腎不全	1						1		1		
(%)	(100.0)						(100.0)				
腎・尿路結核											
(%)											
腎・尿路結石	1						1	1	2		
(%)	(100.0)						(100.0)				
腎・尿路腫瘍	1	1					2	1	3	0.50	0.71
(%)	(50.0)	(50.0)					(100.0)				
閉塞性尿路障害	1						1	2	3		
(%)	(100.0)						(100.0)				
骨髄腫								1	1		
(%)								(100.0)			
腎形成不全	2						2	2	4		
(%)	(100.0)						(100.0)				
不明	64	7	1	1			73	77	150	0.16	0.50
(%)	(87.7)	(9.6)	(1.4)	(1.4)			(100.0)				
再導入	9	1					10	5	15	0.10	0.32
(%)	(90.0)	(10.0)					(100.0)				
その他	22						22	18	40		
(%)	(100.0)						(100.0)				
合計	890	85	13	5	2	2	997	751	1,748	0.15	0.55
(%)	(89.3)	(8.5)	(1.3)	(0.5)	(0.2)	(0.2)	(100.0)				
記載なし (%)											
総計	890	85	13	5	2	2	997	751	1,748	0.15	0.55
(%)	(89.3)	(8.5)	(1.3)	(0.5)	(0.2)	(0.2)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

*個々の患者の腹膜炎発症率(回/1患者・年)は、1.0未満の数値が0となるため、「0」と表示しています。

(C)Japanese Society for Dialysis Therapy

2014年調査