

表 4104 2014年中の出口部感染発症率(回/1患者・年) 原疾患別 (PD・HD(F)等併用患者)

原疾患	0	1.0~	2.0~	3.0~	4.0~	5.0~	合計	不明 記載なし	総計	平均	標準偏差
慢性糸球体腎炎	356	59	15	5	2	6	443	284	727	0.33	0.89
(%)	(80.4)	(13.3)	(3.4)	(1.1)	(0.5)	(1.4)	(100.0)				
慢性腎孟腎炎	11						11	3	14		
(%)	(100.0)						(100.0)				
急速進行性糸球体腎炎	2	2	1	1			6	4	10	1.17	1.17
(%)	(33.3)	(33.3)	(16.7)	(16.7)			(100.0)				
妊娠腎/妊娠中毒症	6	1					7	3	10	0.14	0.38
(%)	(85.7)	(14.3)					(100.0)				
その他の分類不能の腎炎	8	1		1		1	11	8	19	0.82	1.66
(%)	(72.7)	(9.1)		(9.1)		(9.1)	(100.0)				
多発性囊胞腎	20		1		1	1	23	14	37	0.51	1.45
(%)	(87.0)		(4.3)		(4.3)	(4.3)	(100.0)				
腎硬化症	62	10	2			2	76	67	143	0.36	1.14
(%)	(81.6)	(13.2)	(2.6)			(2.6)	(100.0)				
悪性高血圧	12	1		1			14	8	22	0.29	0.83
(%)	(85.7)	(7.1)		(7.1)			(100.0)				
糖尿病性腎症	213	43	10	3	2	5	276	254	530	0.41	1.07
(%)	(77.2)	(15.6)	(3.6)	(1.1)	(0.7)	(1.8)	(100.0)				
SLE腎炎	4						4	4	8		
(%)	(100.0)						(100.0)				
アミロイド腎	1						1	1	2		
(%)	(100.0)						(100.0)				
痛風腎	5	1					6	1	7	0.17	0.41
(%)	(83.3)	(16.7)					(100.0)				
先天性代謝異常にによる腎不全	1						1		1		
(%)	(100.0)						(100.0)				
腎・尿路結核											
(%)											
腎・尿路結石	1						1	1	2		
(%)	(100.0)						(100.0)				
腎・尿路腫瘍	2						2	1	3		
(%)	(100.0)						(100.0)				
閉塞性尿路障害	1						1	2	3		
(%)	(100.0)						(100.0)				
骨髄腫								1	1		
(%)											
腎形成不全	1	1					2	2	4	0.50	0.71
(%)	(50.0)	(50.0)					(100.0)				
不明	60	4	4	3	1		72	78	150	0.35	0.87
(%)	(83.3)	(5.6)	(5.6)	(4.2)	(1.4)		(100.0)				
再導入	9	1					10	5	15	0.10	0.32
(%)	(90.0)	(10.0)					(100.0)				
その他	18	1			2	1	22	18	40	0.82	2.17
(%)	(81.8)	(4.5)			(9.1)	(4.5)	(100.0)				
合計	793	125	33	14	8	16	989	759	1,748	0.37	1.02
(%)	(80.2)	(12.6)	(3.3)	(1.4)	(0.8)	(1.6)	(100.0)				
記載なし (%)											
総計	793	125	33	14	8	16	989	759	1,748	0.37	1.02
(%)	(80.2)	(12.6)	(3.3)	(1.4)	(0.8)	(1.6)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

個々の患者の出口部感染発症率(回/1患者・年)は、1.0未満の数値が0となるため、「0」と表示しています。

(C)Japanese Society for Dialysis Therapy

2014年調査