

表 4125 PET Cr D/P比 Body Mass Index別（腹膜透析患者）

Body Mass Index	0.5未満	0.5～	0.65～	0.81～	合計	記載なし	総計	平均	標準偏差
14未満		4	7	2	13	14	27	0.68	0.10
(%)		(30.8)	(53.8)	(15.4)	(100.0)				
14～	9	10	8	9	36	35	71	0.64	0.19
(%)	(25.0)	(27.8)	(22.2)	(25.0)	(100.0)				
16～	14	39	36	18	107	121	228	0.66	0.14
(%)	(13.1)	(36.4)	(33.6)	(16.8)	(100.0)				
18～	36	113	121	40	310	317	627	0.66	0.14
(%)	(11.6)	(36.5)	(39.0)	(12.9)	(100.0)				
20～	44	199	222	69	534	606	1,140	0.66	0.14
(%)	(8.2)	(37.3)	(41.6)	(12.9)	(100.0)				
22～	55	215	244	89	603	658	1,261	0.66	0.14
(%)	(9.1)	(35.7)	(40.5)	(14.8)	(100.0)				
24～	29	163	215	62	469	439	908	0.67	0.12
(%)	(6.2)	(34.8)	(45.8)	(13.2)	(100.0)				
26～	25	98	120	45	288	273	561	0.67	0.14
(%)	(8.7)	(34.0)	(41.7)	(15.6)	(100.0)				
28～	22	54	54	14	144	158	302	0.63	0.14
(%)	(15.3)	(37.5)	(37.5)	(9.7)	(100.0)				
30～	15	44	45	7	111	109	220	0.64	0.12
(%)	(13.5)	(39.6)	(40.5)	(6.3)	(100.0)				
合計	249	939	1,072	355	2,615	2,730	5,345	0.66	0.13
(%)	(9.5)	(35.9)	(41.0)	(13.6)	(100.0)				
記載なし	63	199	168	53	483	3,113	3,596	0.62	0.16
(%)	(13.0)	(41.2)	(34.8)	(11.0)	(100.0)				
総計	312	1,138	1,240	408	3,098	5,843	8,941	0.65	0.14
(%)	(10.1)	(36.7)	(40.0)	(13.2)	(100.0)				
平均	23.12	23.23	23.35	22.91	23.22	23.07	23.15		
標準偏差	4.11	3.63	3.50	3.42	3.60	3.60	3.60		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査