

表 4130 PET Cr D/P比 補正カルシウム濃度(mg/dL)別 (腹膜透析患者)

| 補正<br>カルシウム濃度 | 0.5未満  | 0.5～   | 0.65～  | 0.81～  | 合計      | 記載なし  | 総計    | 平均   | 標準偏差 |
|---------------|--------|--------|--------|--------|---------|-------|-------|------|------|
| 7.0未満         | 2      | 1      | 3      |        | 6       | 23    | 29    | 0.62 | 0.13 |
| (%)           | (33.3) | (16.7) | (50.0) |        | (100.0) |       |       |      |      |
| 7.0～          |        | 6      | 11     | 2      | 19      | 17    | 36    | 0.68 | 0.10 |
| (%)           |        | (31.6) | (57.9) | (10.5) | (100.0) |       |       |      |      |
| 7.5～          | 6      | 13     | 18     | 4      | 41      | 86    | 127   | 0.64 | 0.16 |
| (%)           | (14.6) | (31.7) | (43.9) | (9.8)  | (100.0) |       |       |      |      |
| 8.0～          | 14     | 48     | 48     | 9      | 119     | 167   | 286   | 0.64 | 0.12 |
| (%)           | (11.8) | (40.3) | (40.3) | (7.6)  | (100.0) |       |       |      |      |
| 8.4～          | 2      | 11     | 15     | 4      | 32      | 54    | 86    | 0.67 | 0.13 |
| (%)           | (6.3)  | (34.4) | (46.9) | (12.5) | (100.0) |       |       |      |      |
| 8.5～          | 48     | 173    | 176    | 42     | 439     | 467   | 906   | 0.64 | 0.14 |
| (%)           | (10.9) | (39.4) | (40.1) | (9.6)  | (100.0) |       |       |      |      |
| 9.0～          | 67     | 285    | 316    | 115    | 783     | 819   | 1,602 | 0.66 | 0.14 |
| (%)           | (8.6)  | (36.4) | (40.4) | (14.7) | (100.0) |       |       |      |      |
| 9.5～          | 87     | 277    | 308    | 102    | 774     | 855   | 1,629 | 0.66 | 0.13 |
| (%)           | (11.2) | (35.8) | (39.8) | (13.2) | (100.0) |       |       |      |      |
| 10.0～         | 9      | 35     | 38     | 10     | 92      | 99    | 191   | 0.66 | 0.14 |
| (%)           | (9.8)  | (38.0) | (41.3) | (10.9) | (100.0) |       |       |      |      |
| 10.1～         | 18     | 88     | 99     | 61     | 266     | 291   | 557   | 0.69 | 0.14 |
| (%)           | (6.8)  | (33.1) | (37.2) | (22.9) | (100.0) |       |       |      |      |
| 10.5～         | 13     | 36     | 55     | 20     | 124     | 172   | 296   | 0.68 | 0.13 |
| (%)           | (10.5) | (29.0) | (44.4) | (16.1) | (100.0) |       |       |      |      |
| 11.0～         | 4      | 12     | 16     | 5      | 37      | 74    | 111   | 0.66 | 0.13 |
| (%)           | (10.8) | (32.4) | (43.2) | (13.5) | (100.0) |       |       |      |      |
| 11.5～         |        | 2      | 7      | 2      | 11      | 18    | 29    | 0.71 | 0.09 |
| (%)           |        | (18.2) | (63.6) | (18.2) | (100.0) |       |       |      |      |
| 12.0～         |        | 1      | 3      | 2      | 6       | 21    | 27    | 0.75 | 0.15 |
| (%)           |        | (16.7) | (50.0) | (33.3) | (100.0) |       |       |      |      |
| 合計            | 270    | 988    | 1,113  | 378    | 2,749   | 3,163 | 5,912 | 0.66 | 0.14 |
| (%)           | (9.8)  | (35.9) | (40.5) | (13.8) | (100.0) |       |       |      |      |
| 記載なし          | 42     | 150    | 127    | 30     | 349     | 2,680 | 3,029 | 0.62 | 0.15 |
| (%)           | (12.0) | (43.0) | (36.4) | (8.6)  | (100.0) |       |       |      |      |
| 総計            | 312    | 1,138  | 1,240  | 408    | 3,098   | 5,843 | 8,941 | 0.65 | 0.14 |
| (%)           | (10.1) | (36.7) | (40.0) | (13.2) | (100.0) |       |       |      |      |
| 平均            | 9.35   | 9.37   | 9.40   | 9.55   | 9.41    | 9.41  | 9.41  |      |      |
| 標準偏差          | 0.78   | 0.69   | 0.75   | 0.70   | 0.73    | 0.85  | 0.80  |      |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査