

表 4131 PET Cr D/P比 リン濃度(mg/dL)別 (腹膜透析患者)

リン濃度	0.5未満	0.5～	0.65～	0.81～	合計	記載なし	総計	平均	標準偏差
3未満	5	17	21	6	49	77	126	0.65	0.15
(%)	(10.2)	(34.7)	(42.9)	(12.2)	(100.0)				
3.0～	13	34	49	12	108	162	270	0.66	0.12
(%)	(12.0)	(31.5)	(45.4)	(11.1)	(100.0)				
3.5～	28	86	87	34	235	290	525	0.65	0.15
(%)	(11.9)	(36.6)	(37.0)	(14.5)	(100.0)				
4.0～	26	138	148	53	365	405	770	0.66	0.13
(%)	(7.1)	(37.8)	(40.5)	(14.5)	(100.0)				
4.5～	37	171	187	56	451	470	921	0.66	0.12
(%)	(8.2)	(37.9)	(41.5)	(12.4)	(100.0)				
5.0～	51	149	199	68	467	481	948	0.66	0.15
(%)	(10.9)	(31.9)	(42.6)	(14.6)	(100.0)				
5.5～	34	162	156	52	404	435	839	0.66	0.13
(%)	(8.4)	(40.1)	(38.6)	(12.9)	(100.0)				
6.0～	13	19	23	9	64	74	138	0.63	0.18
(%)	(20.3)	(29.7)	(35.9)	(14.1)	(100.0)				
6.1～	17	75	76	31	199	278	477	0.67	0.13
(%)	(8.5)	(37.7)	(38.2)	(15.6)	(100.0)				
6.5～	21	60	73	24	178	221	399	0.65	0.14
(%)	(11.8)	(33.7)	(41.0)	(13.5)	(100.0)				
7.0～	9	36	53	15	113	99	212	0.67	0.13
(%)	(8.0)	(31.9)	(46.9)	(13.3)	(100.0)				
7.5～	4	21	24	10	59	90	149	0.68	0.12
(%)	(6.8)	(35.6)	(40.7)	(16.9)	(100.0)				
8.0～	5	15	8	6	34	37	71	0.64	0.15
(%)	(14.7)	(44.1)	(23.5)	(17.6)	(100.0)				
8.5～		6	5	1	12	26	38	0.67	0.10
(%)		(50.0)	(41.7)	(8.3)	(100.0)				
9.0～	8	10	9	4	31	58	89	0.61	0.15
(%)	(25.8)	(32.3)	(29.0)	(12.9)	(100.0)				
合計	271	999	1,118	381	2,769	3,203	5,972	0.66	0.14
(%)	(9.8)	(36.1)	(40.4)	(13.8)	(100.0)				
記載なし	41	139	122	27	329	2,640	2,969	0.62	0.15
(%)	(12.5)	(42.2)	(37.1)	(8.2)	(100.0)				
総計	312	1,138	1,240	408	3,098	5,843	8,941	0.65	0.14
(%)	(10.1)	(36.7)	(40.0)	(13.2)	(100.0)				
平均	5.32	5.23	5.21	5.26	5.23	5.26	5.25		
標準偏差	1.43	1.26	1.26	1.29	1.28	1.41	1.35		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査