

表 4137 PET Cr D/P比 Non-HDL-C濃度(mg/dL)別 (腹膜透析患者)

Non-HDL-C濃度	0.5未満	0.5～	0.65～	0.81～	合計	記載なし	総計	平均	標準偏差
50未満		4	8	2	14	4	18	0.70	0.10
(%)		(28.6)	(57.1)	(14.3)	(100.0)				
50～	3	22	24	10	59	55	114	0.69	0.13
(%)	(5.1)	(37.3)	(40.7)	(16.9)	(100.0)				
70～	17	59	73	27	176	175	351	0.66	0.15
(%)	(9.7)	(33.5)	(41.5)	(15.3)	(100.0)				
90～	36	114	130	55	335	325	660	0.66	0.14
(%)	(10.7)	(34.0)	(38.8)	(16.4)	(100.0)				
110～	39	144	168	56	407	390	797	0.66	0.14
(%)	(9.6)	(35.4)	(41.3)	(13.8)	(100.0)				
130～	31	122	147	47	347	315	662	0.66	0.14
(%)	(8.9)	(35.2)	(42.4)	(13.5)	(100.0)				
150～	29	87	84	34	234	202	436	0.65	0.15
(%)	(12.4)	(37.2)	(35.9)	(14.5)	(100.0)				
170～	18	53	53	15	139	124	263	0.63	0.14
(%)	(12.9)	(38.1)	(38.1)	(10.8)	(100.0)				
190～	5	19	26	3	53	67	120	0.65	0.11
(%)	(9.4)	(35.8)	(49.1)	(5.7)	(100.0)				
210～	6	13	21	4	44	46	90	0.65	0.15
(%)	(13.6)	(29.5)	(47.7)	(9.1)	(100.0)				
合計	184	637	734	253	1,808	1,703	3,511	0.66	0.14
(%)	(10.2)	(35.2)	(40.6)	(14.0)	(100.0)				
記載なし	128	501	506	155	1,290	4,140	5,430	0.65	0.14
(%)	(9.9)	(38.8)	(39.2)	(12.0)	(100.0)				
総計	312	1,138	1,240	408	3,098	5,843	8,941	0.65	0.14
(%)	(10.1)	(36.7)	(40.0)	(13.2)	(100.0)				
平均	132.51	128.99	128.29	123.55	128.30	128.65	128.47		
標準偏差	36.37	37.76	39.54	33.85	37.88	38.16	38.01		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査