

表 4217 残腎Kt/V 体重(kg)別（腹膜透析患者）

体重	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
20未満	4			2	1		1		8	30	38	0.68	0.84
(%)	(50.0)			(25.0)	(12.5)		(12.5)		(100.0)				
20～	3	1	2				2		8	15	23	0.68	0.88
(%)	(37.5)	(12.5)	(25.0)				(25.0)		(100.0)				
30～	10	11	10	9	2		1	1	44	128	172	0.52	0.60
(%)	(22.7)	(25.0)	(22.7)	(20.5)	(4.5)		(2.3)	(2.3)	(100.0)				
40～	74	75	64	45	20	7	5	6	296	608	904	0.55	0.67
(%)	(25.0)	(25.3)	(21.6)	(15.2)	(6.8)	(2.4)	(1.7)	(2.0)	(100.0)				
50～	126	119	115	78	48	11	6	6	509	1,082	1,591	0.55	0.64
(%)	(24.8)	(23.4)	(22.6)	(15.3)	(9.4)	(2.2)	(1.2)	(1.2)	(100.0)				
60～	123	121	138	96	53	18	8	7	564	1,052	1,616	0.60	0.67
(%)	(21.8)	(21.5)	(24.5)	(17.0)	(9.4)	(3.2)	(1.4)	(1.2)	(100.0)				
70～	60	62	70	47	26	3	3	3	274	537	811	0.57	0.68
(%)	(21.9)	(22.6)	(25.5)	(17.2)	(9.5)	(1.1)	(1.1)	(1.1)	(100.0)				
80～	26	21	22	19	10	6	1		105	193	298	0.58	0.55
(%)	(24.8)	(20.0)	(21.0)	(18.1)	(9.5)	(5.7)	(1.0)		(100.0)				
90～	9	1	6	9	8			1	34	70	104	0.74	0.61
(%)	(26.5)	(2.9)	(17.6)	(26.5)	(23.5)			(2.9)	(100.0)				
100～	3		1						4	16	20	0.13	0.25
(%)	(75.0)		(25.0)						(100.0)				
110～		2	1						3	7	10	0.40	0.26
(%)		(66.7)	(33.3)						(100.0)				
120～										9	9		
(%)													
合計	438	413	429	305	168	45	27	24	1,849	3,747	5,596	0.57	0.65
(%)	(23.7)	(22.3)	(23.2)	(16.5)	(9.1)	(2.4)	(1.5)	(1.3)	(100.0)				
記載なし	72	56	80	44	27	11	12	5	307	3,038	3,345	0.63	0.71
(%)	(23.5)	(18.2)	(26.1)	(14.3)	(8.8)	(3.6)	(3.9)	(1.6)	(100.0)				
総計	510	469	509	349	195	56	39	29	2,156	6,785	8,941	0.58	0.66
(%)	(23.7)	(21.8)	(23.6)	(16.2)	(9.0)	(2.6)	(1.8)	(1.3)	(100.0)				
平均	60.69	60.18	61.48	61.81	63.11	63.52	55.37	59.35	61.14	60.55	60.74		
標準偏差	14.21	12.09	12.77	13.06	13.33	11.38	16.82	14.40	13.16	14.24	13.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査