

表 4218 残腎Kt/V Body Mass Index別（腹膜透析患者）

Body Mass Index	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
14未満	2		1						3	24	27	0.13	0.23
(%)	(66.7)		(33.3)						(100.0)				
14～	9	3	3	5			2	1	23	48	71	0.60	0.82
(%)	(39.1)	(13.0)	(13.0)	(21.7)			(8.7)	(4.3)	(100.0)				
16～	11	14	19	11	5		2	2	64	164	228	0.62	0.65
(%)	(17.2)	(21.9)	(29.7)	(17.2)	(7.8)		(3.1)	(3.1)	(100.0)				
18～	52	49	44	35	21	1	1	3	206	421	627	0.53	0.56
(%)	(25.2)	(23.8)	(21.4)	(17.0)	(10.2)	(0.5)	(0.5)	(1.5)	(100.0)				
20～	101	90	90	61	23	5	3	4	377	763	1,140	0.49	0.56
(%)	(26.8)	(23.9)	(23.9)	(16.2)	(6.1)	(1.3)	(0.8)	(1.1)	(100.0)				
22～	84	105	93	63	38	16	7	6	412	849	1,261	0.61	0.70
(%)	(20.4)	(25.5)	(22.6)	(15.3)	(9.2)	(3.9)	(1.7)	(1.5)	(100.0)				
24～	71	73	77	58	30	9	5	3	326	582	908	0.60	0.71
(%)	(21.8)	(22.4)	(23.6)	(17.8)	(9.2)	(2.8)	(1.5)	(0.9)	(100.0)				
26～	32	46	44	29	19	8	4	2	184	377	561	0.65	0.77
(%)	(17.4)	(25.0)	(23.9)	(15.8)	(10.3)	(4.3)	(2.2)	(1.1)	(100.0)				
28～	24	15	30	19	12	3	1	1	105	197	302	0.60	0.55
(%)	(22.9)	(14.3)	(28.6)	(18.1)	(11.4)	(2.9)	(1.0)	(1.0)	(100.0)				
30～	23	9	20	14	13	2	1	1	83	137	220	0.62	0.58
(%)	(27.7)	(10.8)	(24.1)	(16.9)	(15.7)	(2.4)	(1.2)	(1.2)	(100.0)				
合計	409	404	421	295	161	44	26	23	1,783	3,562	5,345	0.57	0.65
(%)	(22.9)	(22.7)	(23.6)	(16.5)	(9.0)	(2.5)	(1.5)	(1.3)	(100.0)				
記載なし	101	65	88	54	34	12	13	6	373	3,223	3,596	0.61	0.69
(%)	(27.1)	(17.4)	(23.6)	(14.5)	(9.1)	(3.2)	(3.5)	(1.6)	(100.0)				
総計	510	469	509	349	195	56	39	29	2,156	6,785	8,941	0.58	0.66
(%)	(23.7)	(21.8)	(23.6)	(16.2)	(9.0)	(2.6)	(1.8)	(1.3)	(100.0)				
平均	23.08	23.06	23.47	23.31	24.01	24.54	23.08	22.60	23.32	23.06	23.15		
標準偏差	3.69	3.18	3.58	3.56	3.70	2.79	4.37	4.01	3.54	3.62	3.60		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査