

表 4222 残腎Kt/V CRP濃度(mg/dL)別（腹膜透析患者）

CRP濃度	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
0.2未満	268	255	257	182	98	24	12	16	1,112	1,955	3,067	0.56	0.67
(%)	(24.1)	(22.9)	(23.1)	(16.4)	(8.8)	(2.2)	(1.1)	(1.4)	(100.0)				
0.2～	95	66	83	60	31	8	5	3	351	644	995	0.56	0.67
(%)	(27.1)	(18.8)	(23.6)	(17.1)	(8.8)	(2.3)	(1.4)	(0.9)	(100.0)				
0.5～	36	31	26	20	10	1	7	2	133	316	449	0.54	0.63
(%)	(27.1)	(23.3)	(19.5)	(15.0)	(7.5)	(0.8)	(5.3)	(1.5)	(100.0)				
1.0～	17	20	17	9	6	3			72	236	308	0.50	0.51
(%)	(23.6)	(27.8)	(23.6)	(12.5)	(8.3)	(4.2)			(100.0)				
2.0～	13	12	10	14	5	1	1		56	146	202	0.56	0.53
(%)	(23.2)	(21.4)	(17.9)	(25.0)	(8.9)	(1.8)	(1.8)		(100.0)				
4.0～	6	3	6	2	2		1		20	63	83	0.54	0.58
(%)	(30.0)	(15.0)	(30.0)	(10.0)	(10.0)		(5.0)		(100.0)				
6.0～	1	6	1			1			9	31	40	0.36	0.53
(%)	(11.1)	(66.7)	(11.1)			(11.1)			(100.0)				
8.0～			1						1	23	24	0.70	
(%)			(100.0)						(100.0)				
10.0～	6	2	2	2	2			1	15	19	34	0.61	0.92
(%)	(40.0)	(13.3)	(13.3)	(13.3)	(13.3)			(6.7)	(100.0)				
15.0～	1	1	2						4	3	7	0.33	0.30
(%)	(25.0)	(25.0)	(50.0)						(100.0)				
合計	443	396	405	289	154	38	26	22	1,773	3,436	5,209	0.56	0.65
(%)	(25.0)	(22.3)	(22.8)	(16.3)	(8.7)	(2.1)	(1.5)	(1.2)	(100.0)				
記載なし	67	73	104	60	41	18	13	7	383	3,349	3,732	0.69	0.68
(%)	(17.5)	(19.1)	(27.2)	(15.7)	(10.7)	(4.7)	(3.4)	(1.8)	(100.0)				
総計	510	469	509	349	195	56	39	29	2,156	6,785	8,941	0.58	0.66
(%)	(23.7)	(21.8)	(23.6)	(16.2)	(9.0)	(2.6)	(1.8)	(1.3)	(100.0)				
平均	0.60	0.54	0.56	0.44	0.49	0.53	0.52	0.80	0.54	0.67	0.63		
標準偏差	2.06	1.58	1.94	1.12	1.47	1.35	0.92	2.85	1.74	1.70	1.71		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査