

表 4226 残腎Kt/V PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
20未満	3	3	7	2	1				16	24	40	0.48	0.35
(%)	(18.8)	(18.8)	(43.8)	(12.5)	(6.3)				(100.0)				
20～	5	9	8	4	1				27	18	45	0.43	0.37
(%)	(18.5)	(33.3)	(29.6)	(14.8)	(3.7)				(100.0)				
35～	10	8	6	1					25	30	55	0.22	0.27
(%)	(40.0)	(32.0)	(24.0)	(4.0)					(100.0)				
50～	7	11	13	6			2		39	38	77	0.49	0.52
(%)	(17.9)	(28.2)	(33.3)	(15.4)			(5.1)		(100.0)				
75～	13	12	8	2	2				37	32	69	0.30	0.35
(%)	(35.1)	(32.4)	(21.6)	(5.4)	(5.4)				(100.0)				
100～	7	5	17	8		2			39	32	71	0.53	0.42
(%)	(17.9)	(12.8)	(43.6)	(20.5)		(5.1)			(100.0)				
125～	7	6	7	6	2				28	17	45	0.45	0.42
(%)	(25.0)	(21.4)	(25.0)	(21.4)	(7.1)				(100.0)				
150～				1					1		1	0.90	
(%)				(100.0)					(100.0)				
151～	6	4	6	2	4			1	23	20	43	0.63	0.80
(%)	(26.1)	(17.4)	(26.1)	(8.7)	(17.4)			(4.3)	(100.0)				
180～	2	4	5	1					12	22	34	0.35	0.32
(%)	(16.7)	(33.3)	(41.7)	(8.3)					(100.0)				
210～	4	2	10						16	11	27	0.39	0.28
(%)	(25.0)	(12.5)	(62.5)						(100.0)				
240～	1	5	1	3					10	13	23	0.49	0.37
(%)	(10.0)	(50.0)	(10.0)	(30.0)					(100.0)				
300～		2	3	3					8	11	19	0.66	0.32
(%)		(25.0)	(37.5)	(37.5)					(100.0)				
350～	2		1	1					4	4	8	0.35	0.44
(%)	(50.0)		(25.0)	(25.0)					(100.0)				
400～	2	3	1						6	5	11	0.20	0.23
(%)	(33.3)	(50.0)	(16.7)						(100.0)				
500～	1	2	1	2					6	5	11	0.42	0.40
(%)	(16.7)	(33.3)	(16.7)	(33.3)					(100.0)				
合計	70	76	94	42	10	2	2	1	297	282	579	0.44	0.44
(%)	(23.6)	(25.6)	(31.6)	(14.1)	(3.4)	(0.7)	(0.7)	(0.3)	(100.0)				
記載なし	2	9	17	4	4		1	1	38	102	140	0.65	0.62
(%)	(5.3)	(23.7)	(44.7)	(10.5)	(10.5)		(2.6)	(2.6)	(100.0)				
総計	72	85	111	46	14	2	3	2	335	384	719	0.46	0.47
(%)	(21.5)	(25.4)	(33.1)	(13.7)	(4.2)	(0.6)	(0.9)	(0.6)	(100.0)				
平均	125.30	132.78	124.86	153.19	113.80	109.00	52.50	166.00	130.17	135.07	132.56		
標準偏差	126.79	129.49	99.65	136.84	59.55	12.73	2.12		118.18	140.41	129.39		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査