

表 4274 残腎Kt/V 原疾患別 (PD・HD(F)等併用患者)

原疾患	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	113	25	8	8	7	3	1		165	562	727	0.21	0.45
(%)	(68.5)	(15.2)	(4.8)	(4.8)	(4.2)	(1.8)	(0.6)		(100.0)				
慢性腎盂腎炎	6	1		1					8	6	14	0.15	0.39
(%)	(75.0)	(12.5)		(12.5)					(100.0)				
急速進行性糸球 体腎炎	2								2	8	10		
(%)	(100.0)								(100.0)				
妊娠腎/妊娠中毒 症	4								4	6	10		
(%)	(100.0)								(100.0)				
その他の分類不能 の腎炎	3						1		4	15	19	0.57	1.15
(%)	(75.0)						(25.0)		(100.0)				
多発性嚢胞腎	5	2		1	1				9	28	37	0.29	0.53
(%)	(55.6)	(22.2)		(11.1)	(11.1)				(100.0)				
腎硬化症	23	10	2	1	2	1		1	40	103	143	0.25	0.56
(%)	(57.5)	(25.0)	(5.0)	(2.5)	(5.0)	(2.5)		(2.5)	(100.0)				
悪性高血圧	5	2							7	15	22	0.04	0.08
(%)	(71.4)	(28.6)							(100.0)				
糖尿病性腎症	52	31	12	2	4	2			103	427	530	0.22	0.39
(%)	(50.5)	(30.1)	(11.7)	(1.9)	(3.9)	(1.9)			(100.0)				
SLE腎炎	1								1	7	8		
(%)	(100.0)								(100.0)				
アミロイド腎										2	2		
(%)													
痛風腎										7	7		
(%)													
先天性代謝異常に よる腎不全	1								1		1		
(%)	(100.0)								(100.0)				
腎・尿路結核													
(%)													
腎・尿路結石										2	2		
(%)													
腎・尿路腫瘍										3	3		
(%)													
閉塞性尿路障害	1								1	2	3		
(%)	(100.0)								(100.0)				
骨髄腫										1	1		
(%)													
腎形成不全										4	4		
(%)													
不明	25	2	1	2				2	32	118	150	0.28	0.78
(%)	(78.1)	(6.3)	(3.1)	(6.3)				(6.3)	(100.0)				
再導入	4								4	11	15		
(%)	(100.0)								(100.0)				
その他	4	1							5	35	40	0.02	0.04
(%)	(80.0)	(20.0)							(100.0)				
合計	249	74	23	15	14	6	2	3	386	1,362	1,748	0.21	0.48
(%)	(64.5)	(19.2)	(6.0)	(3.9)	(3.6)	(1.6)	(0.5)	(0.8)	(100.0)				
記載なし													
(%)													
総計	249	74	23	15	14	6	2	3	386	1,362	1,748	0.21	0.48
(%)	(64.5)	(19.2)	(6.0)	(3.9)	(3.6)	(1.6)	(0.5)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査