

表 4287 残腎Kt/V 腹膜炎罹患回数(回/年)別 (PD・HD(F)等併用患者)

| 腹膜炎罹患回数 | 0.1未満 | 0.1～ | 0.4～ | 0.8～ | 1.2～ | 1.7～ | 2.0～ | 2.4～ | 合計 | 記載なし | 総計 | 平均 | 標準偏差 |
|---------|---------|--------|--------|--------|---------|-------|-------|---------|---------|-------|-------|------|------|
| なし | 227 | 61 | 17 | 13 | 11 | 6 | 2 | 2 | 339 | 551 | 890 | 0.20 | 0.46 |
| (%) | (67.0) | (18.0) | (5.0) | (3.8) | (3.2) | (1.8) | (0.6) | (0.6) | (100.0) | | | | |
| 1回 | 16 | 4 | 3 | 1 | 1 | | | | 25 | 62 | 87 | 0.18 | 0.37 |
| (%) | (64.0) | (16.0) | (12.0) | (4.0) | (4.0) | | | | (100.0) | | | | |
| 2回 | 1 | | 1 | 1 | | | | | 3 | 9 | 12 | 0.47 | 0.50 |
| (%) | (33.3) | | (33.3) | (33.3) | | | | | (100.0) | | | | |
| 3回 | | | | | | | | 1 | 1 | 4 | 5 | 3.50 | |
| (%) | | | | | | | | (100.0) | (100.0) | | | | |
| 4回 | | | | | 1 | | | | 1 | | 1 | 1.30 | |
| (%) | | | | | (100.0) | | | | (100.0) | | | | |
| 5回 | 1 | | | | | | | | 1 | | 1 | | |
| (%) | (100.0) | | | | | | | | (100.0) | | | | |
| 6回 | | | | | | | | | | | | | |
| (%) | | | | | | | | | | | | | |
| 7回 | | | | | | | | | | | | | |
| (%) | | | | | | | | | | | | | |
| 8回 | | | | | | | | | | | | | |
| (%) | | | | | | | | | | | | | |
| 9回以上 | | | | | | | | | | 1 | 1 | | |
| (%) | | | | | | | | | | | | | |
| 合計 | 245 | 65 | 21 | 15 | 13 | 6 | 2 | 3 | 370 | 627 | 997 | 0.21 | 0.49 |
| (%) | (66.2) | (17.6) | (5.7) | (4.1) | (3.5) | (1.6) | (0.5) | (0.8) | (100.0) | | | | |
| 不明 | 2 | 8 | 2 | | 1 | | | | 13 | 6 | 19 | 0.27 | 0.31 |
| (%) | (15.4) | (61.5) | (15.4) | | (7.7) | | | | (100.0) | | | | |
| 記載なし | 2 | 1 | | | | | | | 3 | 729 | 732 | 0.10 | 0.17 |
| (%) | (66.7) | (33.3) | | | | | | | (100.0) | | | | |
| 総計 | 249 | 74 | 23 | 15 | 14 | 6 | 2 | 3 | 386 | 1,362 | 1,748 | 0.21 | 0.48 |
| (%) | (64.5) | (19.2) | (6.0) | (3.9) | (3.6) | (1.6) | (0.5) | (0.8) | (100.0) | | | | |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査