

表 4354 PD Kt/V 腹膜炎罹患回数(回/年)別 (腹膜透析単独患者)

腹膜炎罹患回数	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
なし	8	94	185	360	603	225	120	73	1,668	2,090	3,758	1.33	0.79
(%)	(0.5)	(5.6)	(11.1)	(21.6)	(36.2)	(13.5)	(7.2)	(4.4)	(100.0)				
1回	1	15	20	46	58	16	13	7	176	264	440	1.26	0.70
(%)	(0.6)	(8.5)	(11.4)	(26.1)	(33.0)	(9.1)	(7.4)	(4.0)	(100.0)				
2回	1	1	3	3	12	4	4		28	42	70	1.35	0.56
(%)	(3.6)	(3.6)	(10.7)	(10.7)	(42.9)	(14.3)	(14.3)		(100.0)				
3回		2			1		1		4	18	22	0.95	0.97
(%)		(50.0)			(25.0)		(25.0)		(100.0)				
4回			3		1				4	2	6	0.82	0.39
(%)			(75.0)		(25.0)				(100.0)				
5回				1		1			2	1	3	1.40	0.71
(%)				(50.0)		(50.0)			(100.0)				
6回										1	1		
(%)													
7回													
(%)													
8回										1	1		
(%)													
9回以上													
(%)													
合計	10	112	211	410	675	246	138	80	1,882	2,419	4,301	1.32	0.78
(%)	(0.5)	(6.0)	(11.2)	(21.8)	(35.9)	(13.1)	(7.3)	(4.3)	(100.0)				
不明			13	10	8	1			32	28	60	0.97	0.36
(%)			(40.6)	(31.3)	(25.0)	(3.1)			(100.0)				
記載なし		4	2	9	10	2	2		29	2,798	2,827	1.11	0.54
(%)		(13.8)	(6.9)	(31.0)	(34.5)	(6.9)	(6.9)		(100.0)				
総計	10	116	226	429	693	249	140	80	1,943	5,245	7,188	1.31	0.77
(%)	(0.5)	(6.0)	(11.6)	(22.1)	(35.7)	(12.8)	(7.2)	(4.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査