

表 4420 総透析量 腹膜炎罹患回数(回/年)別 (腹膜透析単独患者)

腹膜炎罹患回数	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	2.8～	合計	記載なし	総計	平均	標準偏差
なし	6	26	39	60	358	402	331	133	119	1,474	2,284	3,758	1.93	0.99
(%)	(0.4)	(1.8)	(2.6)	(4.1)	(24.3)	(27.3)	(22.5)	(9.0)	(8.1)	(100.0)				
1回	1	6	3	5	41	50	28	14	13	161	279	440	1.91	0.99
(%)	(0.6)	(3.7)	(1.9)	(3.1)	(25.5)	(31.1)	(17.4)	(8.7)	(8.1)	(100.0)				
2回	1		1	2	2	5	6	1	5	23	47	70	1.97	0.86
(%)	(4.3)		(4.3)	(8.7)	(8.7)	(21.7)	(26.1)	(4.3)	(21.7)	(100.0)				
3回			1				2			3	19	22	1.63	0.90
(%)			(33.3)				(66.7)			(100.0)				
4回				2			1	1		4	2	6	1.75	0.89
(%)				(50.0)			(25.0)	(25.0)		(100.0)				
5回							1	1		2	1	3	2.45	0.35
(%)							(50.0)	(50.0)		(100.0)				
6回											1	1		
(%)														
7回														
(%)														
8回											1	1		
(%)														
9回以上														
(%)														
合計	8	32	44	69	401	457	369	150	137	1,667	2,634	4,301	1.93	0.99
(%)	(0.5)	(1.9)	(2.6)	(4.1)	(24.1)	(27.4)	(22.1)	(9.0)	(8.2)	(100.0)				
不明				1	9	12	4	2	4	32	28	60	1.93	0.53
(%)				(3.1)	(28.1)	(37.5)	(12.5)	(6.3)	(12.5)	(100.0)				
記載なし		2	1	2	7	4	5		1	22	2,805	2,827	1.58	0.65
(%)		(9.1)	(4.5)	(9.1)	(31.8)	(18.2)	(22.7)		(4.5)	(100.0)				
総計	8	34	45	72	417	473	378	152	142	1,721	5,467	7,188	1.93	0.98
(%)	(0.5)	(2.0)	(2.6)	(4.2)	(24.2)	(27.5)	(22.0)	(8.8)	(8.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査