

表 4427 腹膜炎罹患回数(回/年) 透析歴(年)別 (腹膜透析患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,904	190	28	9	3	1	1		1		2,137	24	1,439	3,600
(%)	(89.1)	(8.9)	(1.3)	(0.4)	(0.1)	(0.0)	(0.0)		(0.0)		(100.0)			
2年～	1,616	213	31	9	3	3					1,875	49	1,277	3,201
(%)	(86.2)	(11.4)	(1.7)	(0.5)	(0.2)	(0.2)					(100.0)			
5年～	874	103	16	7	1					1	1,002	6	649	1,657
(%)	(87.2)	(10.3)	(1.6)	(0.7)	(0.1)					(0.1)	(100.0)			
10年～	177	18	4	2							201		139	340
(%)	(88.1)	(9.0)	(2.0)	(1.0)							(100.0)			
15年～	55		3								58		39	97
(%)	(94.8)		(5.2)								(100.0)			
20年～	10	2									12		8	20
(%)	(83.3)	(16.7)									(100.0)			
25年～	7										7		1	8
(%)	(100.0)										(100.0)			
30年～	5	1									6		4	10
(%)	(83.3)	(16.7)									(100.0)			
35年～	3										3			3
(%)	(100.0)										(100.0)			
40年～														
(%)														
合計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,556	8,936
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
不明													5	5
(%)														
記載なし														
(%)														
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	3.10	3.11	3.68	3.52	2.14	1.75			1.00	5.00	3.11	2.41	3.10	3.10
標準偏差	3.67	3.21	3.76	3.61	2.19	0.50					3.63	1.65	3.50	3.56

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査