

表 4440 腹膜炎罹患回数(回/年) 体重(kg)別 (腹膜透析患者)

体重	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
20未満	28	5									33		5	38
(%)	(84.8)	(15.2)									(100.0)			
20～	18	2			1						21		2	23
(%)	(85.7)	(9.5)			(4.8)						(100.0)			
30～	127	13	2						1		143	1	28	172
(%)	(88.8)	(9.1)	(1.4)						(0.7)		(100.0)			
40～	648	78	14	6							746	1	157	904
(%)	(86.9)	(10.5)	(1.9)	(0.8)							(100.0)			
50～	1,116	141	27	6	2	1	1				1,294	10	287	1,591
(%)	(86.2)	(10.9)	(2.1)	(0.5)	(0.2)	(0.1)	(0.1)				(100.0)			
60～	1,183	132	11	9	1						1,336	14	266	1,616
(%)	(88.5)	(9.9)	(0.8)	(0.7)	(0.1)						(100.0)			
70～	575	67	9	3	1	2				1	658	9	144	811
(%)	(87.4)	(10.2)	(1.4)	(0.5)	(0.2)	(0.3)				(0.2)	(100.0)			
80～	220	14	3		2						239	1	58	298
(%)	(92.1)	(5.9)	(1.3)		(0.8)						(100.0)			
90～	68	8	2	1		1					80	1	23	104
(%)	(85.0)	(10.0)	(2.5)	(1.3)		(1.3)					(100.0)			
100～	16	2									18		2	20
(%)	(88.9)	(11.1)									(100.0)			
110～	7	1									8		2	10
(%)	(87.5)	(12.5)									(100.0)			
120～	6	1									7		2	9
(%)	(85.7)	(14.3)									(100.0)			
合計	4,012	464	68	25	7	4	1		1	1	4,583	37	976	5,596
(%)	(87.5)	(10.1)	(1.5)	(0.5)	(0.2)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	639	63	14	2							718	42	2,585	3,345
(%)	(89.0)	(8.8)	(1.9)	(0.3)							(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	60.72	59.92	59.62	60.84	62.04	73.30	57.10		38.70	76.80	60.64	64.79	61.09	60.74
標準偏差	14.01	13.67	12.81	12.28	20.17	16.80					13.97	11.08	13.64	13.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査