

表 4441 腹膜炎罹患回数(回/年) Body Mass Index別（腹膜透析患者）

Body Mass Index	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
14未満	22	1									23		4	27
(%)	(95.7)	(4.3)									(100.0)			
14～	53	8									61	1	9	71
(%)	(86.9)	(13.1)									(100.0)			
16～	165	17	7	1	1				1		192		36	228
(%)	(85.9)	(8.9)	(3.6)	(0.5)	(0.5)				(0.5)		(100.0)			
18～	470	48	7	5	1						531	1	95	627
(%)	(88.5)	(9.0)	(1.3)	(0.9)	(0.2)						(100.0)			
20～	802	97	13	3	1		1				917	5	218	1,140
(%)	(87.5)	(10.6)	(1.4)	(0.3)	(0.1)		(0.1)				(100.0)			
22～	879	105	19	6		1					1,010	11	240	1,261
(%)	(87.0)	(10.4)	(1.9)	(0.6)		(0.1)					(100.0)			
24～	666	88	6	6	2	1					769	9	130	908
(%)	(86.6)	(11.4)	(0.8)	(0.8)	(0.3)	(0.1)					(100.0)			
26～	418	42	8	2	1	1				1	473	2	86	561
(%)	(88.4)	(8.9)	(1.7)	(0.4)	(0.2)	(0.2)				(0.2)	(100.0)			
28～	217	22	3	2	1						245	3	54	302
(%)	(88.6)	(9.0)	(1.2)	(0.8)	(0.4)						(100.0)			
30～	158	17	5			1					181	2	37	220
(%)	(87.3)	(9.4)	(2.8)			(0.6)					(100.0)			
合計	3,850	445	68	25	7	4	1		1	1	4,402	34	909	5,345
(%)	(87.5)	(10.1)	(1.5)	(0.6)	(0.2)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	801	82	14	2							899	45	2,652	3,596
(%)	(89.1)	(9.1)	(1.6)	(0.2)							(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	23.15	23.13	23.26	23.05	23.58	26.13	21.76		17.67	26.57	23.15	23.99	23.08	23.15
標準偏差	3.63	3.45	4.20	3.40	4.26	3.18					3.62	3.32	3.48	3.60

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査