

表 4442 腹膜炎罹患回数(回/年) BUN(mg/dL)別 (腹膜透析患者)

BUN	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
20未満	13										13	1	15	29
(%)	(100.0)										(100.0)			
20～	119	13	6								138	1	44	183
(%)	(86.2)	(9.4)	(4.3)								(100.0)			
30～	452	52	13	5	2	1					525	6	141	672
(%)	(86.1)	(9.9)	(2.5)	(1.0)	(0.4)	(0.2)					(100.0)			
40～	990	121	25	6	1	1	1		1		1,146	6	240	1,392
(%)	(86.4)	(10.6)	(2.2)	(0.5)	(0.1)	(0.1)	(0.1)		(0.1)		(100.0)			
50～	1,172	114	15	4	2	1					1,308	12	293	1,613
(%)	(89.6)	(8.7)	(1.1)	(0.3)	(0.2)	(0.1)					(100.0)			
60～	821	96	4	5						1	927	6	217	1,150
(%)	(88.6)	(10.4)	(0.4)	(0.5)						(0.1)	(100.0)			
70～	397	40	3	3	1	1					445	7	97	549
(%)	(89.2)	(9.0)	(0.7)	(0.7)	(0.2)	(0.2)					(100.0)			
80～	165	23	4	1	1						194	1	50	245
(%)	(85.1)	(11.9)	(2.1)	(0.5)	(0.5)						(100.0)			
90～	57	4	1								62	2	25	89
(%)	(91.9)	(6.5)	(1.6)								(100.0)			
100～	23	3	1								27	1	12	40
(%)	(85.2)	(11.1)	(3.7)								(100.0)			
110～	3	1									4		3	7
(%)	(75.0)	(25.0)									(100.0)			
120～	5	1									6		1	7
(%)	(83.3)	(16.7)									(100.0)			
合計	4,217	468	72	24	7	4	1		1	1	4,795	43	1,138	5,976
(%)	(87.9)	(9.8)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	434	59	10	3							506	36	2,423	2,965
(%)	(85.8)	(11.7)	(2.0)	(0.6)							(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	54.97	54.94	49.44	54.13	56.71	53.25	41.00		46.00	65.00	54.87	56.47	54.97	54.90
標準偏差	15.25	15.23	16.35	15.48	18.90	16.32					15.27	19.87	17.82	15.82

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査