

表 4445 腹膜炎罹患回数(回/年) CRP濃度(mg/dL)別(腹膜透析患者)

CRP濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
0.2未満	2,235	215	30	9	3				1		2,493	15	559	3,067
(%)	(89.7)	(8.6)	(1.2)	(0.4)	(0.1)				(0.0)		(100.0)			
0.2～	718	68	13	4	2					1	806	18	171	995
(%)	(89.1)	(8.4)	(1.6)	(0.5)	(0.2)					(0.1)	(100.0)			
0.5～	308	43	1	2							354	7	88	449
(%)	(87.0)	(12.1)	(0.3)	(0.6)							(100.0)			
1.0～	196	32	5	2		1					236	2	70	308
(%)	(83.1)	(13.6)	(2.1)	(0.8)		(0.4)					(100.0)			
2.0～	124	23	7	1	1		1				157	1	44	202
(%)	(79.0)	(14.6)	(4.5)	(0.6)	(0.6)		(0.6)				(100.0)			
4.0～	45	11	4								60	1	22	83
(%)	(75.0)	(18.3)	(6.7)								(100.0)			
6.0～	25	3	2								30		10	40
(%)	(83.3)	(10.0)	(6.7)								(100.0)			
8.0～	12	2		2							16		8	24
(%)	(75.0)	(12.5)		(12.5)							(100.0)			
10.0～	20	5	2	2							29		5	34
(%)	(69.0)	(17.2)	(6.9)	(6.9)							(100.0)			
15.0～	5	1									6		1	7
(%)	(83.3)	(16.7)									(100.0)			
合計	3,688	403	64	22	6	1	1		1	1	4,187	44	978	5,209
(%)	(88.1)	(9.6)	(1.5)	(0.5)	(0.1)	(0.0)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	963	124	18	5	1	3					1,114	35	2,583	3,732
(%)	(86.4)	(11.1)	(1.6)	(0.4)	(0.1)	(0.3)					(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	0.55	0.88	1.44	2.41	0.54	1.46	3.55		0.06	0.48	0.61	0.46	0.72	0.63
標準偏差	1.56	2.56	2.54	4.47	0.78						1.73	0.82	1.65	1.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査