

表 4447 腹膜炎罹患回数(回/年) リン濃度(mg/dL)別 (腹膜透析患者)

リン濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
3未満	72	9	7								88	1	37	126
(%)	(81.8)	(10.2)	(8.0)								(100.0)			
3.0～	179	32	5	1							217	1	52	270
(%)	(82.5)	(14.7)	(2.3)	(0.5)							(100.0)			
3.5～	365	42	5	1	1						414	6	105	525
(%)	(88.2)	(10.1)	(1.2)	(0.2)	(0.2)						(100.0)			
4.0～	552	50	13	2			1		1		619	4	147	770
(%)	(89.2)	(8.1)	(2.1)	(0.3)			(0.2)		(0.2)		(100.0)			
4.5～	663	79	11	4	3	2					762	4	155	921
(%)	(87.0)	(10.4)	(1.4)	(0.5)	(0.4)	(0.3)					(100.0)			
5.0～	686	66	9	12	1	1					775	3	170	948
(%)	(88.5)	(8.5)	(1.2)	(1.5)	(0.1)	(0.1)					(100.0)			
5.5～	603	72	3	1	2	1					682	5	152	839
(%)	(88.4)	(10.6)	(0.4)	(0.1)	(0.3)	(0.1)					(100.0)			
6.0～	100	11									111	1	26	138
(%)	(90.1)	(9.9)									(100.0)			
6.1～	310	35	5							1	351	6	120	477
(%)	(88.3)	(10.0)	(1.4)							(0.3)	(100.0)			
6.5～	288	21	9	1							319	1	79	399
(%)	(90.3)	(6.6)	(2.8)	(0.3)							(100.0)			
7.0～	155	18	4	2							179	2	31	212
(%)	(86.6)	(10.1)	(2.2)	(1.1)							(100.0)			
7.5～	101	14	1								116	6	27	149
(%)	(87.1)	(12.1)	(0.9)								(100.0)			
8.0～	50	5									55	2	14	71
(%)	(90.9)	(9.1)									(100.0)			
8.5～	18	9									27		11	38
(%)	(66.7)	(33.3)									(100.0)			
9.0～	65	6									71	1	17	89
(%)	(91.5)	(8.5)									(100.0)			
合計	4,207	469	72	24	7	4	1		1	1	4,786	43	1,143	5,972
(%)	(87.9)	(9.8)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	444	58	10	3							515	36	2,418	2,969
(%)	(86.2)	(11.3)	(1.9)	(0.6)							(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	5.25	5.24	4.85	5.15	4.91	4.97	4.40		4.40	6.40	5.25	5.63	5.24	5.25
標準偏差	1.32	1.40	1.43	0.95	0.79	0.43					1.33	1.67	1.41	1.35

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査