

表 4451 腹膜炎罹患回数(回/年) 総コレステロール濃度(mg/dL)別 (腹膜透析患者)

総コレステロール濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
100未満	31	6	1	1							39		11	50
(%)	(79.5)	(15.4)	(2.6)	(2.6)							(100.0)			
100～	116	10	4								130	2	28	160
(%)	(89.2)	(7.7)	(3.1)								(100.0)			
120～	317	33	8	4	1	1					364		84	448
(%)	(87.1)	(9.1)	(2.2)	(1.1)	(0.3)	(0.3)					(100.0)			
140～	599	57	13	2	1						672	6	128	806
(%)	(89.1)	(8.5)	(1.9)	(0.3)	(0.1)						(100.0)			
160～	647	73	14	5	1		1				741	9	115	865
(%)	(87.3)	(9.9)	(1.9)	(0.7)	(0.1)		(0.1)				(100.0)			
180～	648	74	6	1		1					730	6	97	833
(%)	(88.8)	(10.1)	(0.8)	(0.1)		(0.1)					(100.0)			
200～	421	53	4	1							479	4	81	564
(%)	(87.9)	(11.1)	(0.8)	(0.2)							(100.0)			
220～	249	25	6								280	1	37	318
(%)	(88.9)	(8.9)	(2.1)								(100.0)			
240～	122	12	1	1							136		23	159
(%)	(89.7)	(8.8)	(0.7)	(0.7)							(100.0)			
260～	111	15	1	1							128	2	19	149
(%)	(86.7)	(11.7)	(0.8)	(0.8)							(100.0)			
合計	3,261	358	58	16	3	2	1				3,699	30	623	4,352
(%)	(88.2)	(9.7)	(1.6)	(0.4)	(0.1)	(0.1)	(0.0)				(100.0)			
記載なし	1,390	169	24	11	4	2			1	1	1,602	49	2,938	4,589
(%)	(86.8)	(10.5)	(1.5)	(0.7)	(0.2)	(0.1)			(0.1)	(0.1)	(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	180.04	181.12	167.17	168.00	154.67	154.00	161.00				179.85	179.13	174.05	179.01
標準偏差	40.54	39.56	39.60	44.71	19.22	36.77					40.46	38.48	40.77	40.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査