

表 4452 腹膜炎罹患回数(回/年) HDL-C濃度(mg/dL)別(腹膜透析患者)

HDL-C濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
20未満	4	1		1							6		2	8
(%)	(66.7)	(16.7)		(16.7)							(100.0)			
20～	205	27	6	1							239	2	47	288
(%)	(85.8)	(11.3)	(2.5)	(0.4)							(100.0)			
30～	684	82	13	5						1	785	6	142	933
(%)	(87.1)	(10.4)	(1.7)	(0.6)						(0.1)	(100.0)			
40～	757	78	12	2	1		1		1		852	4	134	990
(%)	(88.8)	(9.2)	(1.4)	(0.2)	(0.1)		(0.1)		(0.1)		(100.0)			
50～	639	73	10	2	2						726	5	108	839
(%)	(88.0)	(10.1)	(1.4)	(0.3)	(0.3)						(100.0)			
60～	397	43	6								446	2	68	516
(%)	(89.0)	(9.6)	(1.3)								(100.0)			
70～	200	19	4			1					224	1	34	259
(%)	(89.3)	(8.5)	(1.8)			(0.4)					(100.0)			
80～	103	12	1	1							117		24	141
(%)	(88.0)	(10.3)	(0.9)	(0.9)							(100.0)			
90～	47	6	1								54		7	61
(%)	(87.0)	(11.1)	(1.9)								(100.0)			
100～	43	8		1							52		5	57
(%)	(82.7)	(15.4)		(1.9)							(100.0)			
合計	3,079	349	53	13	3	1	1		1	1	3,501	20	571	4,092
(%)	(87.9)	(10.0)	(1.5)	(0.4)	(0.1)	(0.0)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	1,572	178	29	14	4	3					1,800	59	2,990	4,849
(%)	(87.3)	(9.9)	(1.6)	(0.8)	(0.2)	(0.2)					(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	50.75	50.45	48.23	49.92	51.67	76.00	42.00		41.00	39.00	50.68	45.40	49.40	50.47
標準偏差	17.59	19.80	15.87	30.42	5.13						17.84	14.15	17.04	17.72

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査