

表 4453 腹膜炎罹患回数(回/年) Non-HDL-C濃度(mg/dL)別 (腹膜透析患者)

Non-HDL-C濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
50未満	11	3	1								15		3	18
(%)	(73.3)	(20.0)	(6.7)								(100.0)			
50～	81	6	3			1					91		23	114
(%)	(89.0)	(6.6)	(3.3)			(1.1)					(100.0)			
70～	256	25	10	1	1						293		58	351
(%)	(87.4)	(8.5)	(3.4)	(0.3)	(0.3)						(100.0)			
90～	514	48	8	2							572	2	86	660
(%)	(89.9)	(8.4)	(1.4)	(0.3)							(100.0)			
110～	587	75	8	4	1		1				676	7	114	797
(%)	(86.8)	(11.1)	(1.2)	(0.6)	(0.1)		(0.1)				(100.0)			
130～	498	60	9	3							570	1	91	662
(%)	(87.4)	(10.5)	(1.6)	(0.5)							(100.0)			
150～	329	47	4	1							381	3	52	436
(%)	(86.4)	(12.3)	(1.0)	(0.3)							(100.0)			
170～	204	23	1								228	1	34	263
(%)	(89.5)	(10.1)	(0.4)								(100.0)			
190～	93	8	2	1							104	1	15	120
(%)	(89.4)	(7.7)	(1.9)	(1.0)							(100.0)			
210～	70	7	1								78	1	11	90
(%)	(89.7)	(9.0)	(1.3)								(100.0)			
合計	2,643	302	47	12	2	1	1				3,008	16	487	3,511
(%)	(87.9)	(10.0)	(1.6)	(0.4)	(0.1)	(0.0)	(0.0)				(100.0)			
記載なし	2,008	225	35	15	5	3			1	1	2,293	63	3,074	5,430
(%)	(87.6)	(9.8)	(1.5)	(0.7)	(0.2)	(0.1)			(0.0)	(0.0)	(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	128.97	130.50	116.34	130.58	102.00	52.00	119.00				128.89	141.94	125.44	128.47
標準偏差	38.19	35.70	38.29	32.22	19.80						37.96	36.73	38.27	38.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査