

表 4459 腹膜炎罹患回数(回/年) 一日透析液使用量(L/日)別(腹膜透析患者)

一日透析液使用量	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2未満	101	8									109		10	119
(%)	(92.7)	(7.3)									(100.0)			
2～	332	35	5	2	1		1				376	3	14	393
(%)	(88.3)	(9.3)	(1.3)	(0.5)	(0.3)		(0.3)				(100.0)			
4～	911	96	12	4	2				1		1,026		46	1,072
(%)	(88.8)	(9.4)	(1.2)	(0.4)	(0.2)				(0.1)		(100.0)			
6～	1,483	186	34	9	1	3					1,716	2	77	1,795
(%)	(86.4)	(10.8)	(2.0)	(0.5)	(0.1)	(0.2)					(100.0)			
8～	1,171	118	22	6	2	1				1	1,321	17	68	1,406
(%)	(88.6)	(8.9)	(1.7)	(0.5)	(0.2)	(0.1)				(0.1)	(100.0)			
10～	345	41	7	3	1						397	2	11	410
(%)	(86.9)	(10.3)	(1.8)	(0.8)	(0.3)						(100.0)			
12～	174	16		2							192	1	8	201
(%)	(90.6)	(8.3)		(1.0)							(100.0)			
合計	4,517	500	80	26	7	4	1		1	1	5,137	25	234	5,396
(%)	(87.9)	(9.7)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	134	27	2	1							164	54	3,327	3,545
(%)	(81.7)	(16.5)	(1.2)	(0.6)							(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	6.79	6.75	6.77	7.33	6.30	7.18	3.00		5.20	8.00	6.79	7.59	6.61	6.78
標準偏差	2.74	2.63	1.88	3.43	2.93	1.41					2.72	2.34	2.61	2.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査