

表 4463 腹膜炎罹患回数(回/年) 1日のPD実施時間(時間/日)別(腹膜透析患者)

1日のPD実施時間	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
1～	79	6	2		1						88		10	98
(%)	(89.8)	(6.8)	(2.3)		(1.1)						(100.0)			
5～	725	67	11	3	1	1			1		809	1	22	832
(%)	(89.6)	(8.3)	(1.4)	(0.4)	(0.1)	(0.1)			(0.1)		(100.0)			
9～	587	70	8	5		1					671	1	29	701
(%)	(87.5)	(10.4)	(1.2)	(0.7)		(0.1)					(100.0)			
13～	351	37	6		1						395	1	7	403
(%)	(88.9)	(9.4)	(1.5)		(0.3)						(100.0)			
18～	198	12			1						211		2	213
(%)	(93.8)	(5.7)			(0.5)						(100.0)			
24	2,524	305	51	19	3	2	1			1	2,906	1	97	3,004
(%)	(86.9)	(10.5)	(1.8)	(0.7)	(0.1)	(0.1)	(0.0)			(0.0)	(100.0)			
合計	4,464	497	78	27	7	4	1		1	1	5,080	4	167	5,251
(%)	(87.9)	(9.8)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
記載なし	187	30	4								221	75	3,394	3,690
(%)	(84.6)	(13.6)	(1.8)								(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
平均	18.26	18.84	19.01	19.78	16.71	16.25	24.00		8.00	24.00	18.33	14.25	17.75	18.31
標準偏差	7.12	6.95	7.20	6.70	8.58	8.96					7.11	7.41	7.74	7.13

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査