

表 4470 腹膜炎罹患回数(回/年) 年齢(歳)別 (腹膜透析単独患者)

年齢	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
15歳未満	41	8			1						50		47	97
(%)	(82.0)	(16.0)			(2.0)						(100.0)			
15歳～	51	8									59	2	31	92
(%)	(86.4)	(13.6)									(100.0)			
30歳～	293	18	3	1							315	2	208	525
(%)	(93.0)	(5.7)	(1.0)	(0.3)							(100.0)			
45歳～	916	86	14	4		1					1,021	13	645	1,679
(%)	(89.7)	(8.4)	(1.4)	(0.4)		(0.1)					(100.0)			
60歳～	1,706	194	29	9	3	2	1				1,944	29	1,260	3,233
(%)	(87.8)	(10.0)	(1.5)	(0.5)	(0.2)	(0.1)	(0.1)				(100.0)			
75歳～	701	118	22	7	2				1		851	13	596	1,460
(%)	(82.4)	(13.9)	(2.6)	(0.8)	(0.2)				(0.1)		(100.0)			
90歳～	50	8	2	1							61	1	40	102
(%)	(82.0)	(13.1)	(3.3)	(1.6)							(100.0)			
合計	3,758	440	70	22	6	3	1		1		4,301	60	2,827	7,188
(%)	(87.4)	(10.2)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			
不明														
(%)														
記載なし														
(%)														
総計	3,758	440	70	22	6	3	1		1		4,301	60	2,827	7,188
(%)	(87.4)	(10.2)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			
平均	62.83	64.96	67.67	67.55	64.50	66.00	61.00		77.00		63.16	64.20	63.37	63.25
標準偏差	14.78	15.46	12.89	13.12	25.81	7.00					14.85	13.97	15.40	15.06

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査