

表 4476 腹膜炎罹患回数(回/年) 使用透析液の種類別(腹膜透析単独患者)

使用透析液の種類	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
1.5%液のみ	1,745	178	31	9	1	1					1,965	9	117	2,091
(%)	(88.8)	(9.1)	(1.6)	(0.5)	(0.1)	(0.1)					(100.0)			
1.5%液と2.5%液の併用	433	69	12	4	1	1					520	7	38	565
(%)	(83.3)	(13.3)	(2.3)	(0.8)	(0.2)	(0.2)					(100.0)			
2.5%液のみ	98	14	4								116	2	8	126
(%)	(84.5)	(12.1)	(3.4)								(100.0)			
4.25%液のみ(イコデキストリンなし)	4	3									7			7
(%)	(57.1)	(42.9)									(100.0)			
イコデキストリン単独(ブドウ糖使用なし)	47	2									49		2	51
(%)	(95.9)	(4.1)									(100.0)			
1.5%+イコデキストリン	825	96	11	3	1		1		1		938	2	65	1,005
(%)	(88.0)	(10.2)	(1.2)	(0.3)	(0.1)		(0.1)		(0.1)		(100.0)			
1.5%+2.5%+イコデキストリン	305	30	7	2	2						346		20	366
(%)	(88.2)	(8.7)	(2.0)	(0.6)	(0.6)						(100.0)			
2.5%+イコデキストリン	234	33	5	4	1	1					278	3	13	294
(%)	(84.2)	(11.9)	(1.8)	(1.4)	(0.4)	(0.4)					(100.0)			
4.25%液+イコデキストリン	2	1									3			3
(%)	(66.7)	(33.3)									(100.0)			
合計	3,693	426	70	22	6	3	1		1		4,222	23	263	4,508
(%)	(87.5)	(10.1)	(1.7)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			
不明	9										9	37	2	48
(%)	(100.0)										(100.0)			
記載なし	56	14									70		2,562	2,632
(%)	(80.0)	(20.0)									(100.0)			
総計	3,758	440	70	22	6	3	1		1		4,301	60	2,827	7,188
(%)	(87.4)	(10.2)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査